INDEX

COPEIA, 1983, NOS. 1–4

NEW NAMES

Bagarius ***suchus*** 435–437, 439, 441–443
Campostoma ***pauciradii*** 101–116
Ctenoblepharis ***adiutuvelatus*** 454–457, 790
Gambusia ***xanthosoma*** 457–464
Gastrotheca ***dendronastes*** 868–874
Liolaemus ***fabiani*** 788–790
Litoria ***umbonata*** 803–808
Minascaecilia sartoria 857–863
Minyichthys ***inusitatus*** 774–776
Modicus minimus 863–867
Modicus tangaroa 863–867
Odax ***cyanoallix*** 95–101
Oedura ***gemmata*** 445–449
Paracaesio ***stonei*** 450–453
Pseudoeurycea ***longicauda*** 884–894
Rana (*Paa*) ***rara*** 895–901
Toyatamaphimeia 89–95

NEW AND ESTABLISHED NAMES

Ablabes purpurans 799
Abronia 1070–1071
Abudefduf saxatilis 830
Acanthistius 779
Acanthophis 200 *antarcticus* 202
Acanthophora sp. 1019
Acanthurus coeruleus 830
Acer saccharhinum 340
Acheta domestica 835, 838
Acris 156 *crepitans* 300–311
Acropora palmata 831
Adenostoma fasciculatum 511
Aeluroscalabotes 466–467 *felinus* 467
Afroedura 467–473
Agalychnis 641
  callidryas 426, 641–642, 644
  saltator 641–642, 644
Agama agama 937
Agamura 467, 469, 473
Agkistrodon 221–224 *piscivorus* 221
Ailuronyx 467, 473
Albula vulpes 699
Algansea aphanea 501
Alligator 89
  mississippiensis 388, 842–843
  sinensis 93
Allodontichthys 498, 501, 507
  hubbsi 499–501, 507–508
  sp. 499–500, 502
  tamazulae 499–502, 507–508
  zonistius 499–501
Allognathosuchus mooki 844
Alloophorus 498, 501, 503, 509 *robustus* 499–502
Allotoca 498, 501, 509
  dugesi 499–502
  maculata 499–502
Alopias 375
  pelagicus 375
  superciliosus 375–382
  vulpinus 375
Alsodes 30–31, 34–36, 1104–1107
  barrioi 1104, 1106–1107
  gargola 1104–1106
  monticola 1104–1106
  nodosus 1104, 1106–1107
  pehuenche 1105
  tumultuosus 1104, 1106
  vanzolinii 1104–1107
  verrucosus 1104–1107
Alsophylax 467, 469, 475
Ambloplites
  constellatus 1042, 1049
  rupestris 540
Ambrosia dumosa 269–270
Ambystoma 76, 609, 627–629, 1001, 1004
  fluvinatum 68
  gracile 627
  jeffersonianum 1001–1005, 1037–1038, 1040
  mabeii 630
  maculatum 615, 630, 1001–1005
  mexicanum 981
  ordinarium 889
  rosaceum 67–78
  rosaceum nigrum 68
  rosaceum rosaceum 68
  rosaceum sonoraensis 68
  texanum 624–628
  tigrinum 67–78, 608–616, 627–639, 1004, 1006
  tigrinum mavortium 1006–1012
  tigrinum nebulosum 68, 77, 1006–1007, 1011
  tigrinum stebbensi 68
  tigrinum tigrinum 608–616, 628–629
  tigrinum velasci 68
Ameca 498, 505, 507, 509 *splendens* 499–500, 502–503
Ameiva
  ameiva 364
  spp. 744
Amphibolurus
  muricatus 201
  spp. 201
Amphignathodon 869, 872–873 *guentheri* 869, 872–874

Amphiprion
melanopus 712
spp. 812
Amphiuma 627
Anarbylus 465–467
Anarchopterus 774
Anchoa 48, 50, 53
analis 50
arenicola 48–49
argenteus 50
argentivittata 48–54
belizensis 50
cayorum 50
chamensis 50–51
choerostoma 50
clarki 50–51, 53
colonensis 50
compressa 50–51
cubana 50
curta 51–53
delicatissima 50–53
duodecim 48, 50
eigenmannia 50–51, 53
elongata 50
exigua 49–53
filifera 50
ginsburgi 50
helleri 49–51, 53
hepsetus 50
ischana 49–51
januaria 50
lamprotaenia 50
lucida 50, 53
lyolepis 48, 50
marinii 50
mitchilli 50
mundeola 48, 50
mundeoloides 50
naso 50–51
nasus 50–51
nasuta 50
panamensis 48, 50–53
parva 50
pectoralis 50
schultzi 51
scofieldi 50–51, 53
spinifer 50–51
starksi 50–51, 53
tricolor 50
trinitatis 50
tropica 51, 53
ubatubae 50
walkeri 48, 50–53
Anchovia
arenicola 49, 51
clupeoides 50
eigenmannia 51
macrolepidota 50
miarcha 49
mundeoloides 51
nigra 50
scitula 49
surinamensis 50
Anchoviella 50, 53
analis 50–51
balboae 50, 52
blackburni 50
brevirostris 50
carrikeri 50
cayennensis 50
guianensis 50
helleri 49
jamesi 50
lepidentostole 50
miarcha 48–49
nattereri 50
parri 50
perfasciata 49–50
scitula 48–50
vaillanti 50
Ancylodactylus 473
Andrena erythronii 710
Aneides ferreus 667
Anniella 79–89
geronimensis 79, 81–83
nigra 79, 81–85
nigra argentea 79–81, 86–87
nigra nigra 79–80, 84–87
pulchra 79–87
pulchra nigra 79, 82–84
pulchra pulchra 79, 82–83, 86
texana 79, 82, 84–86
Anodus 618
Anolis 131, 139, 320, 430, 523
aeneus 328
carolinensis 158–159, 212, 523–529, 701–704, 836
conspersus 328, 363–364
cristatellus 703
cupreus 329
distichus 328–329
garmani 364
limifrons 420–434
lineatopus 320–322
oculatus 328
opalinus 319–331
polylepis 328
porcatus 523, 527
roquet 328, 701
sagrei 328–329, 433
valencienni 322
Anoplarchus purpurescens 316
Anoplocapros 902, 912–917
amygdaloides 904, 911, 916
grayi 916
lenticularis 916
Anthias 782 *squamipinnis* 812
Anyperodon 784 *leucogrammicus* 778
Apeltes quadracus 541–544
Aporops 784 *bilinearis* 778
Aracana 902, 912–917
aurita 904, 910, 916
ornata 916

Aracanostracion 902
Archoplites 780
Aristelliger 467, 473
Artemia 542, 552
Artemisia 9 *californica* 511
Asaccus 467–468
Ascaphus 689
Aspasma 866
Aspasmichthys 866
Aspasmodes 866
Ataeniobus 498, 503, 505, 507–509 *toweri* 497, 499–500, 502–503
Atelognathus 34–36
Atriplex 9
Augochloropsis diversipennis 710
Aulacocephalus 782, 784 *temmincki* 778
Aulostomus maculatus 548, 830
Australocrinia 275 *riparia* 275
Austrelaps 200

Baccharis pilularis 511
Badis badis 441
Bagarius 435–445
  bagarius 435–444
  Buchanani 437
  lica 435, 438–439
  nieuwenhuisi 435
  Nieuwenhuisi 438
  suchus 435–437, 439, 441–443
  yarrelli 435–441, 443–444
Balsadichthys 505, 508
Basiliscus 228
  basiliscus 477
  plumifrons 477
Batea catharinensis 334
Batrachophrynus 35–36
Batrachoseps 1052, 1067–1068
Batrachus 15
Batrachyla 34–35
Belonoperca 784 *chabanaudi* 778
Betta splendens 975, 977
Blaesodactylus 473
Blechnum blechnoides 33
Bodianus rufus 830
Bogertia 467, 473
Boiga 204
Boleosoma
  lepida 1082
  phlox 1082–1083
Bolitoglossa 1067, 1070–1071 *subpalmata* 1037–1038, 1040–1041
Bombina orientalis 987
Boop boops 814
Bouteloua-Hilaria 9
Brachiodontes exustus 333
Brachydanio rerio 243, 1093
Brachyurabochampsa 93
Briba 467, 473
Bucephalus longicornutus 863
Bufo 420
  americanus 426
  bufo 153
  woodhousei 426
Bunopus 467, 469, 473
Buteo jamaicenis 836

Cacophis 200
Caecilia 857–858, 861–862
  ochrocephala 862
  volcani 859
Caiman 844
  crocodilus 93, 842–843
  crocodilus yacare 92
Callagur borneoensis 923
Callanthias 781
Callendula sp. 992
Callisaurus draconoides 270
Calodactylodes 467, 473
Campostoma 101–102, 104–108, 110–112, 114–115, 1076
  anomalum 102, 106, 538, 824
  anomalum anomalum 111–112
  anomalum michauxi 101–102, 104–107, 110–112, 114–115
  anomalum pullum 101–102, 104–108, 110–112, 114–115
  oligolepis 101–107, 109–115
  oligolepis × *Campostoma pauciradii* 115
  ornatum 102, 111–112
  pauciradii 101–116
Canthigaster rostrata 830
Caprichthys 902, 912–917 *gymnura* 904, 912, 916
Caprichthys-Capropygia lineage 915
Capropygia 902, 912–917 *unistriata* 916
Caranx bartholomaei 251
Carassius auratus 116–121, 746, 753
Carcharhinus
  dussumieri 496
  longimanus 544
Carcharodon 126 *carcharias* 126
Caretta 516, 518 *caretta* 516–519, 924, 1013, 1020
Carettochelys insculpta 925–926
Carya glabra 340
Cassidix mexicanus 309
Castor canadensis 478
Catostomus 44
  clarki 47, 316
  commersoni 63, 246–250, 540
  discobolus 37–39, 41–47
  discobolus discobolus 38
  discobolus yarrowi 37–48
  platyrhynchus 47
  plebius 37–39, 41–47
  santaanae 47
  yarrowi 37, 43–44, 46
Caudiverbera 35
Centrarchops 781
Centropristis striata 783
Cephalanthus occidentalis 609
Cepphus columba 511

Cetengraulis
edentulus 50
juruensis 50
mysticetus 50
Cetorhinus 126 *maximus* 126
Chaenogaleus 487 *macrostoma* 487
Chaetodon 828
capistratus 830
ocellatus 830
Chanda siamensis 441
Chaoborus 545
Chapalichthys 498, 501, 503, 509
encaustus 499–500, 502–503
pardalis 499–500, 502–503
peraticus 503
Characodon 498, 503, 507 *lateralis* 499–500, 503–504
Chasmatosaurus 485
Cheilinus 875, 877
fasciatus 875, 877–878, 881
nigropinnatus 876, 879, 881
Chelonia 123, 841, 926
mydas 121–125, 215, 218, 839–842, 924, 928, 1013–1023
mydas agassizii 1021
Chelydra
mydas 382
serpentina 152, 215, 218, 264–268, 384, 386, 388, 394, 414, 926
serpentina acutirostris 924
serpentina osceola 924
serpentina serpentina 924, 1086
Chinemys reevesi 920, 922
Chirolophius
laticeps 13
malabaricus 13–14
Chironius exoletus 799
Chironomus plumosus 605
Chiropterotriton 1071
Chondrodactylus 467–468, 473, 477
Chondropoulos & Lykakis 991–1001
Chorizanthe spp. 269–270
Chriopeops 564 *goodei* 564, 569
Chromis
cyaneus 830
multilineatus 830
Chrosomus
erythrogaster 825
neogaeus 822–826
Chrysanthemum sp. 992
Chrysemys
concinna 921
concinna mobilensis 921
concinna texana 921
decorata 921
floridana 921
floridana hoyi 921
malonei 921
nelsoni 921
picta 214–221, 264–268, 347–358, 921
picta belli 348–349, 353–355, 522–523, 921
picta dorsalis 348–349, 354–355, 921
picta marginata 357, 921
picta picta 1086
rubiventris 921
scripta callirostris 921
(*Pseudemys*) *scripta elegans* 921
scripta ornata 921
stejnegeri vicina 921
terrapen 921
Chrysophrys auratus 316
Chusquea quila 33
Claudius angustatus 923, 925
Clemmys
guttata 922
insculpta 922
marmorata marmorata 922
marmorata pallida 922
muhlenbergi 922
Clepticus parrai 830
Clinocottus
analis 695
populations 695
Cnemaspis 467, 469, 473
Cnemidophorus 211, 359, 362, 364, 991, 997
deppii 364
flagellicaudus 586
gularis 156
hyperythrus 211–212
inornatus 211
lemniscatus 359, 364
neomexicanus 211
ocellifer 359–366
sexlineatus 206–214
sp. 996, 998
tigris 212, 270, 586–587, 997–998
trigris multiscutatus 996
Coelorhynchus carminatus carminatus 1084–1085
Coilia 53
Coldenia palmeri 269–270
Coleodactylus 466
amazonicus 466
brachystoma 466
meridionalis 466
Coleonyx 465–467
brevis 465
elegans 465
mitratus 465
reticulatus 465
variegatus 465–466
Colopus 467–468, 475
Colossoma 618
Coluber 221, 937
bicolor 795, 800
constrictor 211, 512, 937
dictyodes 795, 800
merremii 795, 800
miliaris 793–794, 799
Constrictor 221–224 *constrictor* 221–222
Copeotyphlinus 857, 862 *syntremus* 859
Cophosaurus texanus 586–587
Coronella
australis 795, 800

orientalis 799
poecilolaemus 796, 800
Corophium sp. 334
Corvus brachyrynchous 511
Coryphaenoides
alateralis 1083–1086
rupestris 1084–1085
thelestomus 1083–1086
Coryphopterus
nicholsi 809–812
personatus 809, 830
Cosymbotus 467, 473
Cottus carolinae 975–977
Couesius plumbeus 825
Crenichthys 497, 507
Crocodylus 89, 93–94, 844
acutus 90, 93, 388, 390, 844
intermedius 90
niloticus 93, 388, 390
porosus 93
porosus biporcatus 92
rhombifer 93
"*Crocodylus*" *cataphractus* 93
"*Crocodylus* sp." 844
Cromileptes 784 *altivelis* 778
Crossobamon 467, 469, 473
Crossocheilus 441
Crotalus 221–224, 262, 520
durissus 519–522
durissus terrificus 519–522
viridis 211–213, 221, 519–522, 586
viridis viridis 519–522
Crotaphytus 1–11, 478
bicinctores 1, 3, 5, 9–10
collaris 6, 9, 477, 481, 484, 586, 835–837
collaris baileyi 1–11
collaris bicinctores 9
dickersonae 10
insularis 6, 10
insularis bicinctores 1–11
insularis vestigium 9–10
wislizenii 268, 270
Cryophis 1070
Cryptantha spp. 269–270
Cryptobalis[sic]es 914
Cryptobranchus 627
alleganiensis 225
alleganiensis alleganiensis 225–226, 228, 230
alleganiensis bishopi 225–231
Ctenoblepharis 454
adspersus 454
audituvelatus 454–457, 790
nigriceps 454
reichei 454, 456
stolzmanni 454
Cypselurus 738
Cyrtocara
annectens 544–546
lateristriga 546
moorii 544–547
rostratus 546
Ctenopharyngodon idella 749–755
Ctenosaura similis 705
Ctenotus taeniolatus 201
Cuora
amboinensis 922
flavomarginata 923
Cyclemys dentata 922
Cyclothone 554, 562
Cymadusa sp. 334–335
Cymothoa exigua 813–815
Cyprinella 1074, 1076
Cyprinodon 499, 507
hubbsi 569
variegatus 569
Cyprinus carpio 116–121, 745–749, 753
Cyrtodactylus 467–469, 472–473, 475
agamuroides 475
agusanensis 475
amictopholis 475
angularis 475
annulatus 475
arnouxii 468–469, 475
baluensis 475
biordinis 475
brevipalmatus 475
brevipes 475
caspius 475
cavernicolus 475
chitralensis 475
collegalensis 475
condorensis 475
consobrinoides 475
consobrinus 475
darmandvillei 475
deccanensis 475
derongo 475
deveti 475
elongatus 475
fasciolatus 475
feae 475
fedtschenkoi 475
fraenatus 475
fumosus 475
galgajuga 468–469, 475
gastropholis 475
gubernatoris 475
heterocercus 475
himalayanus 475
intermedius 475
irregularis 475
jellesmae 475
jeyporensis 475
kachhensis 475
khasiensis 475
kirmanensis 475
kotschyi 475
laevigatus 475
lateralis 475
lawderanus 475
loriae 475
louisiadensis 475

malayanus 475
malcomsmithi 475
marmoratus 475
mimikanus 475
montiumsalsorum 475
nebulosus 475
novaeguineae 475
oldhami 475
papuensis 475
peguensis 475
pelagicus 468–470, 475
pelagicus undulatus 469
philippinicus 475
pubisulcus 475
pulchellus 469, 475
quadrivirgatus 475
redimiculus 475
rubidus 475
russowii 475
sagittifer 475
scaber 475
sermowaiensis 475
serpensinsula 468–469, 475
stoliczkai 475
sworderi 475
triedrus 475
turcmenicus 475
vankampeni 468–469, 475
variegatus 475
watson 475
wetariensis 475
yakhuna 475

Dalea spp. 269–270
Danio malabaricus 975, 977
Dascyllus trimaculatus 1087
Daucus carota 992
Deirochelys
reticularia 922
reticularia chrysea 922
Dellichthys morelandi 863
Delma torquatus 468
Dendrelaphis 204
Dendroica discolor 512
Dermatemys mawii 924, 926, 928
Dermatolepis dermatolepis 544
Dermochelys coriacea 928
Desmognathus 1052, 1068
auriculatus 340
fuscus 1102–1104
fuscus fuscus 1101–1104
monticola 1102–1104
ochrophaeus 1024–1025, 1033–1034, 1098–1102
spp. 624
Diadophis punctatus 586
Dicamptodon 679–691
aterrimus 679–680, 684–685, 687, 689–690
copei 679–680, 683–689
ensatus 72, 679–680, 683–689
Dicrostonyx torquatus 767

Didelphys virginiana 156
Dinoperca 781 *petersii* 778
Diplanthera wrightii 1019
Diplocynodon 92
Diplodactylus 166–168
ciliaris 166
elderi 166–167
michaelseni 166, 168
spinigerus 161–169
steindachneri 168
vittatus 168
williamsi 167, 201
Diploprion 784 *bifasciatum* 778
Diploria labyrinthiformis 831
Dipsas catesbyi 964
Dipsosaurus dorsalis 268–271
Discolophius 15
Distichlis spicata 719, 724
Dogania subplana 925
Dorosoma cepedianum 621
Dravidogecko 467, 473
Dromicus amazonicus 796, 800
Drosophila 162, 196, 236, 298, 748
melanogaster 295
pseudoobscura 295
Drysdalia 200
coronata 967
coronoides 202
Dyrosaurus 842

Ebenavia 467, 473
Echinocactus polycephalus 9
Echiostoma barbatum 555
Eichhornia 618
Elaphe
guttata 837
obsoleta 512
Elasmopus sp. 334
Eleocharis 609
Eleutherodactylus 941–953
antillensis 942–949
biporcatus groups 859
brittoni 942–949, 951
cochranae 942–949
cooki 942–949
coqui 942–949, 951, 1089–1090
eneidae 942–950
gryllus 942–949
hedricki 942–948
jasperi 941
karlschmidti 942–951
locustus 942–949
portoricensis 942–949, 1089–1090
richmondi 942–950
unicolor 942–950
wightmanae 942–949
Empetrichthys 497, 507
Emydoidea blandingi 264–268, 922
Emys orbicularis 921
Encrasicholina 48–54

devisi 52–53
heterolobus 52–53
punctifer 52–53
purpurea 52–53
Engraulis 48, 53
anchoita 50
argentivittatus 51
(*Stolephorus*) *argentivittatus* 48
clarki 51
eurystole 50
mordax 49–50
panamensis 52–53
ringens 50
Eoplectus 903, 905, 914
Ephedra 9
Epibulus 875, 877 *insidiator* 877
Epinephelus 782, 784
(*Alphestes*) 784
(*Cephalopholis*) 784
cruentatus 778
(*Dermatolepis*) 784
(*Epinephelus*) 784
fasciatus 544
flavolimbatus 778
inermis 778
itajara 778
morio 778
multiguttatus 778
niveatus 778
(*Promicrops*) 784
sp. 783
striatus 830
Eremias 1095 *lugubris* 998
Eretmochelys 841 *imbricata* 271–273, 839–842, 924
Erichthonias sp. 334
Esox
lucius 621
niger 1087
Etheostoma 295, 1047
blennioides 1047
caeruleum 625, 1049
caeruleum caeruleum 1049
cinereum 1047
(*Etheostoma*) 1045, 1049–1050
(*Etheostoma*) *zonale* 1045
euzonum 1042–1050
euzonum erizonum 1042–1049
euzonum euzonum 1042–1049
euzonum euzonum × *Etheostoma erizonum* 1044, 1048
euzonum euzonum-Etheostoma euzonum erizonum 1045–1047 intergrades
juliae 1042, 1049
kanawhae 1047, 1050
lepidum 1082
(*Nanostoma*) 1049
nianguae 1047
osburni 1047–1050
sagitta 1047
spectabile 625, 1082
spectabile fragi 1049
spectabile pulchellum 1082
spectabile uniporum 1049
spp. 625
tetrazonum 1047, 1050
variatum 1047, 1050
zonale 1045, 1049
(*Etheostoma*) *zonale* 1045, 1049
Eublepharis 466
hardwickii 467
macularius 467
Eumeces
inexpectatus 212
laticeps 212
obsoletus 586
Euparkeria 485
Euphorbia
sp. 992
spp. 269–270
Eupomacentrus
partitus 830
planifrons 830
spp. 830
variabilis 830
Eupsophus 34–36
coppingeri 34
nodosus 1104
Eurycea
bislineata 340, 627, 1036–1041
longicauda 340
Eurypharynx pelecanoides 560–563
Evechinus chloroticus 863
Exiloboa 1070
Exogone dispar 334

Flagelloserranus 785
Forsterygion varium 832–834
Fundulus 55, 60, 63, 564, 719, 724, 970–971
adinia 64
adinius 64
cingulatus 569
diaphanus 63–64
grandis 564, 711–718, 737
grandissimus 564
heteroclitus 196, 316, 564, 711–712, 715–717, 719–740
heteroclitus grandis 727
heteroclitus heteroclitus 727, 738
heteroclitus macrolepidotus 727, 738
kansae 55, 58, 61–64
(*Plancterus*) *kansae* 55
majalis 564
notti 60
olivaceus 975, 977
persimilis 564
(*Plancterus*) 55
seminolis 569
similis 564
zebra 55
zebrinus 55–67
(*Plancterus*) *zebrinus* 55
Furina 200

Galesphyris 485
Gambelia
silus 10
wislizenii 10
Gambusia 457, 460–463, 598–600, 605–607, 978
affinis 182–183, 243–245, 953–963
affinis affinis 975, 977
alvarezi 179
aurata 179
beebei 457, 459–463
eurystoma 457, 459–463
hurtadoi 179
luma 457, 459–463, 599, 606–607
pseudopunctata 457, 459–462
punctata 457, 459–463
puncticulata 457
puncticulata yucatana 598–607
rhizophorae 457, 459–463
sexradiata 457, 459–463, 598–607
xanthosoma 457–464
Garthia 473
Garzoniella 468
Gasterosteus
aculeatus 259–261, 541–544
wheatlandi 541, 543
Gastrotheca 868–869, 871–873
angustifrons 868–869, 872–874
bufona 868–869, 872–874
ceratophrys 868–869, 872
cornuta 868–869, 872–874
dendronastes 868–874
nicefori 872
weinlandii 872
Gavialis 92–94 *gangeticus* 93
Geckolepis 467, 473
Geckonia 467–468, 473
Gehyra 467, 473
australis 201
mutilata 470
spp. 201
Geisopristurus 468, 470
Gekko 467, 469, 473 *gecko* 701
Genyonemus lineatus 691–696
Geochelone 920
carbonaria 923
chilensis 923
denticulata 923
elephantopus 384, 386–388, 392, 394, 397, 920, 923
elongata 920, 923
gigantea 923
pardalis 920, 923
Geoemyda spengleri 922
Gephyrosaurus 847 *bridensis* 845–847
Gerrhonotus kingi 586
Geukensia demissa 719, 721–724, 738
Gharialis 94
Girardinichthys 498, 503, 505, 508–509
innominatus 508
multiradiatus 499–500, 503–504, 508
viviparus 499–500, 503, 508
Gobiesox 554
eugrammus 552, 554
maeandricus 551–554
papillifer 552, 554
rhessodon 552–554
strumosus 552, 554
Gobiosoma
genie 830
multifasciatum 809
spp. 548
Gomphosus 826 *varius* 828
Gonatodes 466 *vittatus* 465–467
Gonioplectrus 784 *hispanus* 778
Gonostoma
bathyphilum 562
elongatum 554–557
gracile 554, 556
Goodea 498, 503, 507–509
atripinnis 499–500, 503, 505
gracilis 499–500, 503, 505
luitpoldi 499–500, 503, 505
Gopherus 920
agassizi 923
berlandieri 923
polyphemus 923
Gracila 784 *albomarginata* 778
Gracileria sp. 1019
Gramma loreto 829–832
Grammistes 782, 784 *sexlineatus* 778
Grammistops 784 *ocellatus* 778
Graptemys
barbouri 921
caglei 921
flavimaculata 921
geographica 264–268, 921
kohni 921
nigrinoda 921
oculifera 921
ouachitensis 264–268
pseudogeographica 921
pseudogeographica ouachitensis 921
pseudogeographica sabinensis 921
pulchra 921
versa 921
Grus antigone antigone 388–389
Gymnodactylus 467, 475
Gymnophthalmus
multiscutatus 364, 801
rubicauda 801
Gymnoscyphus 863
Gymnothorax griseus 544

Hadropterus shumardi 1083
Halichoeres garnoti 830
Halodule wrightii 1014, 1019
Heliconia 871
Heloderma suspectum 477–478
Hemanthias peruanus 252–253
Hemidactylus 467, 469, 473 *frenatus* 470
Hemigaleus 487–497 *microstoma* 487–497
Hemilutjanus 781 *macrophthalmus* 778

Hemiphyllodactylus 467, 478 *typus* 470
Hemitheconyx 466 *caudicinctus* 467
Heosemys spinosa 922
Hesperocorixa spp. 234
Hesperoleucus symmetricus 699
Heteronotia 467, 470, 473
Heteroscarus 95
Hieremys annandalei 923
Hildebrandichthys setiger 50
Hipopterigium sp. 33
Hippoglossoides platessoides 621
Holacanthus
  ciliaris 830
  tricolor 830
Holodactylus 466 *africanus* 467
Homonota 467, 470, 473
Homopholis 467–468, 473
Hoplocephalus 198–205
  bitorquatus 198–204
  bungaroides 198–204
  spp. 198
  stephensi 198–204
Howella 780
Hubbsina 498, 503, 505, 508–509 *turneri* 499–500, 504–505, 508–509
Hybopsis
  monacha 558, 1074–1076
  monacha × *Notropis galacturus* 1074–1077
Hydrargira 55 *ornata* 738
Hydrargyra 55 *zebra* 55, 64
Hydrolycus 618
Hydromantes 892
Hydromorphus concolor 859
Hyla 641, 643-644, 803, 950, 981, 1070
  avivoca 275
  bifurca 641–642, 644
  bokermanni 641–643, 646
  brevifrons 641–644, 646
  calcarata 641–642, 644
  ceratophrys 872, 874
  chrysoscelis 238–242, 273–275
  chrysoscelis-Hyla versicolor complex 273, 275
  cinerea 156, 513–516, 950, 981
  ebraccata 641–642, 645–647
  geographica 426–427, 641–642, 644
  granosa 641–642, 646
  gratiosa 232–237, 513–516, 950
  lanciformis 641–642, 644, 646
  leucophyllata 640–642, 644, 646
  loquax 641–642
  marmorata 641–642, 644
  melanogyrea 641–642, 644
  minuscula 641–642, 644–647
  minuta 641–642
  multifasciata 641–642, 644, 646
  parviceps 641–643, 646–647
  parviceps group 643
  phlebodes 641–642, 644–645
  punctata 641–643, 645
  regilla 513–516
  rhodopepla 641–643, 646–647
  rosenbergi 513
  "*rubra*" 645
  sarayacuensis 641–644
  triangulum 641–644, 646–647
  uranochroa group 859
  versicolor 273, 275
Hylorina 35–36
Hymenophyllum
  caudiculatum 33
  plicatum 33
Hypentelium nigricans 540
Hypophthalmichthys molitrix 753
Hypoplectrus 782 spp. 830

Ichthyomyzon
  castaneus 367–374
  gagei 367–374
Ictalurus
  nebulosus 1087
  punctatus 975, 977
Iguana iguana 477–478, 481, 484, 529–530
Ilyodon 498–499, 501, 505
  furcidens 499–500, 505, 508
  sp. 499, 505
  whitei 499–500, 504–505, 508
  xantusi 499–500, 505, 508
Imantodes cenchoa 964
Insuetophrynus 31, 34–36 *acarpicus* 30–47
Inula sp. 992
Ioa vigil 1082–1083
Isurus 126, 375, 377
  oxyrinchus 126–130, 375–378, 380–381
  paucus 126, 375–382
  spp. 375

Jeboehlkia 784–785 *gladifer* 778
Jordanella floridae 569
Juncus repens 608

Kachuga
  dhongoka 923, 926
  smithi 923
  tecta 923
  trivittata 923
Kaokogecko 467–468, 473
Kentnocapros 902, 911–916
  aculeatus 906, 912, 916
  flavofasciatus 916
  rosapinto 912, 916
Kinixys belliana belliana 923
Kinosternon 925–926
  bauri 924
  baurii 926
  flavescens 924
  herrerai 924
  hirtipes 924
  integrum 924
  leucostomum 924
  leucostomum postinguinale 924
  scorpioides 924, 926

scorpioides abaxillare 924
scorpioides carajasensis 924
scorpioides cruentatum 924
scorpioides scorpioides 924
subrubrum 924, 926
subrubrum hippocrepis 924
subrubrum steindachneri 924
Kyphosus analogus 544
Labrus
bergylta 811
ossifagus 811
Lacerta 991
agilis 997
lepida 477–478, 482
viridis 477, 996–997
vivipara 996
Lamna 126, 375 *nasus* 126, 375–376
Lampetra
fluviatilis 367, 372
japonica 372
lamottenii 373
planeri 367, 372–373
Lampropeltis getulus 937
Lamprophlis guichenoti 201
Larrea tridentata 269–270
Larus glaucescens 511
Lavinia exilicauda 699
Leidyosuchus 844
formidabilis 844
wilsoni 844
?*Leidyosuchus* 843
Leiocassis 441
Lembesseia parvianalis 818
Lembos sp. 334–335
Lemna sp. 302
Lepidoblepharis 466
Lepidochelys kempi 384–385, 388, 1013
Lepidodactylus 467 *lugubris* 470
Lepomis 1081
cyanellus 625
gibbosus 336
macrochirus 336, 540, 1086–1088
megalotis 540, 625, 1087
Leptodactylus albilabris 941
Leptolucania ommata 569
Lermichthys 508
Lethrinops praeorbitalis 544–546
Leuciscus prolixus 102
Leuresthes
sardina 716
tenuis 711
Libellula spp. 234
Limia
caymanensis 457
formosa 816, 818
poecilioides 816
Limnodynastes
fletcheri 201
tasmaniensis 201
Limnomedusa 36
Liobranchia 866
Liolaemus 788–790
anomalus group 790
constanzae 456, 790
fabiani 788–790
insolitus 790
nigroroseus 456
paulinae 456
Liophis 791, 796, 799
amazonicus 791–802
australis 793, 797
chrysostomus 791–802
cobella collaris 799
fuscus 793
miliaris 791–802
miliaris amazonicus 792, 800
miliaris chrysostomus 792, 796, 798–800
miliaris merremii 792–793, 795, 797, 799–800
miliaris miliaris 792–793, 796, 798–800
miliaris mossoroensis 792–793, 797–799, 801
miliaris orina 793
miliaris orinus 792–793, 795–796, 798–800
miliaris semiaureus 792–793, 795, 797, 800
mossoroensis 791–802
orientalis 793
orina 793
poecilogyrus 799
poeciliostomus 793
purpurans 791–802
reginae 799
reginae ornata 793, 795, 800
reginae ornatus 797
undulatus auctorum 799
Liopropoma 782, 784 *susumi* 778
Liquidambar styraciflua 340, 609
Liriodendron tulipifera 340
Lissemys punctata 925
Lissonanchus 866
Litoria 803–808
amboinensis 806–807
caerulea 201
congenita 806–807
darlingtoni 806–807
dentata 807
everetti 806–807
peroni 806–807
rothii 806–807
rubella 201, 806–807
tyleri 807
umbonata 803–808
wisselensis 803, 806–807
Lophidius 15
Lophiodes 12, 14, 25, 28
beroe 16
bruchius 29
caulinaris 12
caulinaris species group 28
kempi 18, 26
miacanthus 12
mutilus 12
olivaceus 12
spilurus 12

Lophiomus 11–30
caulinaris 12
indicus 11, 14
laticephalus 11, 23–24
laticeps 11, 14
lithulon 23
longicephalus 13–14, 24
malabaricus 11
miacanthus 12
olivaceus 12
setigerus 11–14, 24–26, 28
upsicephalus 25–26
viviparus 11
Lophiopsis 15
Lophius 11–30
americanus 11, 14–20, 28
budegassa 11, 14–16, 18–21, 24–27
eurypterus 24
gastrophysus 11, 14–16, 18–23, 29
indicus 11, 13
litulon 11, 14–15, 18–21, 23–24
parvipinnis 16
piscatorius 11, 14–16, 18–21, 24–28
setigerus 12–14
sp. 18
upsicephalus 11, 14–15, 18–21, 25–27
vaillanti 11, 14–15, 18–21, 25–26, 29
viviparus 13–14
vomerinus 11, 14–15, 18–21, 25, 27, 29
Lophosoria quadripinnata 33
Loricaria 258, 618
"*cataphracta*" 258
piracicabae 258
simillima 257–259
Loricariichthys 257–258
labialis 257–259
platymetopon 257–259
spixii 258
Lucania 564–570
goodei 564–569
interioris 564
parva 564–569
Luperosaurus 467, 473
Lupinus sp. 511
Lutjanus 851 *guttatus* 813–816
Luzuriaga radicans 33
Lycengraulis
batesii 50
grossidens 50
group 53
limnichthys 50
olidus 50
poeyi 50, 52
simulator 50
Lygodactylus 467–468, 471, 473

Mabuya
buettneri 138
maculilabris 138
Maccullochella 779
Macroclemys temminckii 924, 926
Macrognathus aculeatus 444
Macrorhamphosus scolopax 251
Malaclemys
terrapin 267, 922
terrapin littoralis 922
Malacochersus tornieri 923
Malayemys subtrijuga 923
Martes pennanti 478
Masticophis
bilineatus 586
lateralis 937
Mauremys
caspica leprosa 922
caspica rivulata 922
japonica 922
mutica 922
Mecistops 93
Meinertia oestroides 814
Melanochelys trijuga coronata 922
Melanotaenia nigrans 975, 977
Melichthys niger 255
Melospiza melodia 512
Menidia 564
beryllina 57, 568–569
colei 568
peninsulae 568–569
Microcaecilia 857, 861 *supernumeraria* 861
Microgecko 469, 472–473
Micrognathus 774 (*Minyichthys*) 774
Micropterus 672, 1087
dolomieui 672–679
punctulatus 672
salmoides 193–197, 1087
sp. 114
treculi 672–679
Microscalabotes 468, 473
Microspathodon chrysurus 712, 830
Microtus 511–512 sp. 511
Millotisaurus 467, 473
Minascaecilia 857–863 ***sartoria*** 857–863
Minomus jarrovii 37
Minyichthys 774–776
brachyrhinus 774, 776
inusitatus 774–776
myersi 774, 776
sentus 774–776
Minytrema melanops 540
Modicus 863–868
minimus 863–868
tangaroa 863–868
Modiolus demissus 719
Mogurnda mogurnda 337
Mollienesia 817–818
latipinna 818
petenensis 818
sphenops altissima 819
sphenops macrura 819
sphenops melanistia 820
sphenops petersi 820

sphenops vantynei 819
sulphuraria 820
velifera 818
Monocanthus ciliatus 331–338
Montastrea annularia 831
Moxostoma 501
atripinne 538–541
carinatum 975, 977
duquesnei 540
eurythrurum 540
lachneri 114
sp. 114
Mugil sp. 753
Mupus maculatus 251
Mus musculus 201, 837
Mustela vison 478
Mycteroperca 784 *phenax* 778, 783
Mylossoma 618
Myopus schisticolor 767
Myrica cerifera 609
Myriophyllum pinnatum 634

Nactus 469–470
arnouxii 469–470
galgajuga 469–470
pelagicus 469–470
serpensinsula 469–470
vankampeni 469–470
Naja 221–224 *naja* 221
Narudasia 467–468, 473
Naso 828
Nasturtium officinale 302
Nasua
narica 530
nasua 706
Naucrates ductor 251, 544
Naultinus 466 *elegans* 466
Necturus 627
Neelaps 200
Negogaleus 487
Nemichthys scolopaceus 562
Neoodax 95–96, 100
Neoophorus 498, 501, 505, 509
catarinae 499–500, 505
diazi 499–500, 505
meeki 499–500, 504–505
regalis 505
sp. 497, 500, 504
Neotoca 508 *bilineata* 508
Nerodia 1087 *sipedon* 1095, 1097
Neurospora 748
Nezumia bairdii 1084–1085
Ninia maculata 967
Niphon 777, 779, 781, 784-786 *spinosus* 777-787
Nocomis 112, 1076
biguttatus 824
platyrhynchus × *Notropis chrysocephalus* 1077
Nomeus 250–252 *gronovii* 250–252
Nosema 815
Notechis 200
Nothofagus 1105
Notonecta spp. 234
Notophagus 36
Notophthalmus
viridescens 340, 628–629, 1037–1038
viridescens dorsalis 628–639
Notropis 112, 1076–1077, 1079
amoenus × *Notropis atherinoides* 1077
ardens 538, 625
atherinoides 1077–1081
atherinoides × *Notropis volucellus* 1077–1082
buchanani 1079
callisema 114
callistius 975–977
callitaenia 114
chrysocephalus 538, 1077, 1079
cornutus 625, 824
cornutus × *Notropis rubellus* 1077
cornutus × *Rhinichthys cataractae* 1077
(*Cyprinella*) 825, 1074, 1076
euryzonus 114
galacturus 1074–1076
(*Cyprinella*) *galacturus* 1074
hypsilepis 114
kanawha 1077
lutrensis 825
pilsbryi 1049
rubellus × *Notropis volucellus* 1077
spilopterus 825, 1074
spp. 625
stramineus 1079
venustus 822–826
volucellus 1077–1081
volucellus volucellus 1081
volucellus wickliffi 1077, 1081
whipplei 822–826, 1074
xaenurus 114
zonatus 1049
Noturus
albater 1042
baileyi 558–560
flavater 1042
flavipinnis 558–560
(*Rabida*) 558
Nucras tesselata 998
Nyctimystes 803
Nyssa biflora 609

Ocadia sinensis 923
Odax 95–96, 100
cyanoallix 95–101
pullus 95–96, 100
Odontaspis 375 *taurus* 375, 377, 379–380
Oedipina 1071
Oedura 445–449
castelnaui 445, 449
coggeri 445
gemmata 445–449
lesueuri 445
marmorata 445–449

monilis 445
reticulata 445
rhombifer 445
robusta 445
tryoni 445
Oikopleura spp. 551
Oligocottus maculatus 833
Olisthops 95
Ollentodon 508 *multipunctatus* 508
Ololygon 641, 643–644
boesemani 641, 643–644
boulengeri 641, 643–644, 646
cruentomma 641, 643–644, 646
egleri 641, 643–644
elaeochroa 641, 643-644
garbei 641, 643-644, 646
parkeri 641, 643–646
rostrata group 644
rubra 641, 643–644
rubra group
"*rubra*" 641
Oncorhynchus 563
gorbuscha 549–551
nerka 621–622
tschawytscha 563
tshawytscha 563
Oostethus 774
Opheodrys aestivus 741–745, 967
Opheomorphus
fuscus 795, 800
merremii semiaureus 800
semiaureus 795
Oriitia borneensis 923
Oryzias latipes 311–318
Oscaecilia 857, 861–862 *ochrocephala* 859, 862
Osmerus 813
eperlanus 813
mordax 621, 812–813
Osteolaemus 92–93 *tetraspis* 92–93
Ostracion 913–915
Oxyuranus 200

Pachydactylus 467–468, 473
Paleothyris 485
Palmatogecko 467–468, 473
Panaspis nimbaensis 138
Pantosteus plebius 37
Papuengraulis 53
Paracaesio 450–453
caeruleus 450, 452–453
cantharoides 450, 452–453
gonzalesi 450, 452–453
kusakarii 450–453
pedleyi 450, 452–453
sordidus 450, 452–453
stonei 450–453
tumidus 450, 452–453
xanthurus 450, 452–453
Paracaprella tenuis 334–335
Paradelma orientalis 468
Paragaleus 487
pectoralis 487
tengi 487
Paragehyra 468, 475
Paragobiodon spp. 809, 812
Paralabrax clathratus 691–696
Paranthias 784 *furcifer* 778
Parascaphirhynchus albus 696
Pareas carinatus 967
Parodax 95
Paroedura 467, 471–472
Parvicaecilia 857, 861
Peprilus 397–398, 402, 404–405
alepidotus 397
burti 397–406
paru 397–406
simillimus 398, 405
triacanthus 397–406
Perca flavescens 1087
Percichthys 779
Percilia 779
Percina 1083
caprodes 975, 977
copelandi 1083
nigrofasciata 975, 977
ouchitae 1083
shumardi 1083
sp. 1083
(*Imostoma*) *uranidea* 1083
Perochirus 467, 475
Petromyzon marinus 367, 372–375
Phalacrocorax pelagicus 511
Phelsuma 467, 473
Phenacobius mirabilis 63
Phoenicopterus ruber ruber 388–389
Pholidosaurus ("*Macrorhynchus*") *meyeri* 844
Phoxinus 112
Phrynocephalus 477 *mystaceus* 479
Phyllodactylus 467–469, 471–473, 475
angelensis 475
angustidigitus 475
annulatus 475
ansorgii 475
apricus 475
barringtonensis 475
baurii 475
bordai 475
bugastrolepis 475
clinatus 475
darwini 475
davisi 475
delcampi 475
dixoni 475
duellmani 475
europaeus 471, 475
galapagensis 475
gerrhopygus 475
gilberti 475
guentheri 475
heterurus 475
homolepidurus 475

inequalis 475
inexpectatus 468, 471–472, 475
insularis 475
interandinus 475
johnwrighti 475
julieni 475
kofordi 475
lanei 475
leei 475
lepidopygus 475
lineatus 475
marmoratus 475
martini 475
melanostictus 475
microlepidotus 475
microphyllus 475
muralis 475
palmatus 468, 471–472, 475
palmeus 475
partidus 475
paucituberculatus 475
peringueyi 475
porphyreus 475
pulcher 471, 475
pumilis 475
reissii 475
riebeckii 468, 475
rutteni 475
santacruzensis 475
sentosus 475
siamensis 475
tinklei 475
trachyrhinus 467–468, 475
transversalis 475
tuberculosus 475
unctus 475
ventralis 475
weileri 468, 471–472, 475
wirshingi 475
wolterstorffi 468, 471–472, 475
xanti 475
Phyllomedusa 641, 643
hypocondrialis 641, 643
palliata 641, 643–644
tarsius 641, 643, 646
vaillanti 420–428
Phyllopezus 467, 473
Phyllurus 470
Physalaemus pustulosus 154
Physalia 250–252 *physalis* 250–252
Pikea 784 *longilepis* 778
Pimelodus bagarius 436–437
Pimephales
notatus 538, 540
promelas 625, 822
Pinus
elliottii 609
taeda 609
Pinus-Juniperus 9
Pituophis 221, 223–224 *melanoleucus* 221–222, 586, 937
Plagioscion 618
Plancterus 55 *zebra* 55
Platynereis dumerilii 334
Platynotus semitaeniatus 131–141, 998
Platypoecilus 970–971 *nelsoni* 819
Platysternon 919 *megacephalum* 923, 925
Plectranthias 782
Plectropomus 784 *maculatus* 778
Plethodon 339, 891–892, 1052, 1067–1068
cinereus 344
dorsalis 339
glutinosus 340, 344, 1024–1026, 1032–1034
hoffmani 344
idahoensis 689
jordani 1024–1035, 1037–1038, 1040
vandykei 689
vehiculum 344
websteri 339–346
Podarcis 991
erhardii 992
melisellensis 992
muralis 992–993, 996–997
peloponnesiaca 992
sicula 993
taurica 991–992
taurica ionica 991–1001
taurica taurica 991–992, 996–997
Poecilia 177, 818, 969–973, 978
catemaconis 819–821
chica 819, 821
chisoyensis 819
formosa 816–818, 820–821, 969–974
formosa "A" 970
formosa "B" 970
formosa "C" 970
formosa × *Poecilia latipinna* 817
gilli 818
latipinna 190, 243–245, 816–818, 821, 969–970, 972–973, 975, 977
latipinna complex 817, 821
latipunctata 819, 821
limantouri 819
mexicana 818–821, 969–971, 973
(*Mollienesia*) 817–822
orri 820
petenensis 818, 821
reticulata 170, 177–178, 243–245, 975, 977
sp. 820
sphenops 819–820, 822
sphenops complex 817–819, 821
spp. 969–973
sulphuraria 820–821
thermalis 819
velifera 818, 821
veti-providentiae 820
Poecilichthys
euzonus 1045
euzonus erizonus 1049
spectabilis 1082
Poeciliopsis 177, 287, 293, 296–298, 875
infans 501

lucida 178, 287–299
monacha 296, 298, 766
monacha-lucida 296, 746
occidentalis 961
viriosa 170, 178
Pogonoperca 782–784 *punctata* 778
Polychrus acutirostris 138–139
Polyodon spathula 698
Polyplacapros 902, 911–917 *tyleri* 902–905, 907, 909, 916
Polyprion 781
americanus 778
arcuatus 830
Potamegeton sp. 544
Priapella intermedia 179
Prionace glauca 130, 495
Prionurus punctatus 544
Pristurus 465, 467–469, 472–473
carteri 468, 472
celerrimus 472
crucifer 468, 472
flavipunctatus 468, 472
insignis 472
minimus 472
phillipsii 472
rupestris 468, 472
simonettai 472
sokotranus 468, 472
Proaracana 909, 911
Prochilodus mariae 617–623
Prodiplocynodon 92–93
Profundulus 507
Pseudacris 156
Pseudechis 200
Pseudemys
scripta 215, 218–219, 264–268, 355, 529–530
scripta elegans 522–523
Pseudobranchus 627
Pseudocarcharias kamoharai 375, 380
Pseudoeurycea 884–894, 1052, 1070–1071
altamontana 884–885, 887–893
bellii 889
cephalica 892
cephalica group 892
gadovii group 892
goebeli 892
leprosa 884–886, 888-893
leprosa group 892
longicauda 884–894
orizabensis 893
rex 892
robertsi 884–885, 887–893
smithi 884, 892
unguidentis 884, 892
"*Pseudoeurycea leprosa*" 893
Pseudogekko 467, 473
Pseudogonatodes 466
Pseudogramma 782, 784 *polycantha* 778
Pseudonaja 200
Pseudothecadactylus 471
Pseudotriton
montanus 1026
spp. 624
Ptenopus 467, 473
Pterengraulis atherinoides 50
Ptychadena mascareniensis 979, 984
Ptychozoon 467
Ptyodactylus 467, 473
Pyxidea mouhouti 922

Quendenfeldtia 467–468, 473
Quercus
douglasii 933
nigra 609
rubra 340

Rainfordia 781–782, 784–785 *opercularis* 778
Rana 407, 897
annandalii 899–901
(*Paa*) *annandalii* 895, 899–901
blanfordi 899–901
(*Paa*) *blanfordii* 895, 899–901
catesbeiana 142, 407–410, 534, 1097
chiricahuensis 981
clamitans 141–155, 238–242, 513, 534, 979, 981–985, 988
esculenta 988–989
hecksheri 427
liebigii 898–901
liebigii group 898–900
liebigii supergroup 899
(*Paa*) *maculosa* 900
nigromaculata 989
(*Paa*) 895–901
pipiens 310, 407, 534, 979–989
polunini 899–901
(*Paa*) *polunini* 895, 899–901
pretiosa 1097
rara 898–901
(*Paa*) ***rara*** 895–901
rostandi 899–901
(*Paa*) *rostandi* 895, 899–901
sylvatica 979, 984, 1096–1098
temporaria 989
utricularia 426, 615
Ranidella 275–279, 534
bilingua 534
glauerti 534
insignifera 278, 534
parinsignifera 534
pseudinsignifera 278
riparia 275–278, 534–537
signifera 275–278, 534
tasmaniensis 534
Rhabdofario lacistris 563
Rhadinaea 799
binotata 799
chrysostoma 800
merremi natricoides 795, 800
orina 795, 800
Rheithrodontomys 512

Rheobatrachus silus 806
Rhinichthys 112
Rhinoclemmys 920
areolata 922
funerea 922
pulcherrima 922
punctularia melanosterna 922
punctularia punctularia 922
rubida 922
Rhizoprionodon acutus 496
Rhoptropella 473
Rhoptropus 467–468, 473
Rhynchoedura 466
Ricola 259
Rimicola 552
dimorpha 554
eigenmanni 554
muscarum 552–554
Rineloricaria 259
Rivulus 295, 507
cylindraceus 297
marmoratus 296–297
Rubus 511
Rypticus 782, 784

Sacalia bealei 920, 922
Salmo 563
aguabonita 651
apache 659
clarki 563
clarki bouvieri 659
clarki clarki 659
clarki henshawi 659
clarki lewisi 659
gairdneri 64, 650–662
gilae 659
mykiss 651
sp. 563
Salvadora hexalepis 586
Saurodactylus 468–468, 473
Sauromalus obesus 270
Saurosternon 485
Scaphirhynchus 696–697
albus 696–700
platorynchus 696–700
Scarus 253, 256
croicensis 254
iserti 254
spp. 830
vetula 253–256
Sceloporus 933, 937–939
clarki 586–587
graciosus 139
jarrovi 933, 938–939
magister 586–587
occidentalis 213, 933–936, 938
occidentalis biseriatus 936
occidentalis occidentalis 933–940
olivaceus 433, 933, 938
orcutti 211, 938
poinsetti 268
teapensis 156–157
undulatus 139, 212, 586, 597, 933
undulatus erythrocheilus 933, 938–939
undulatus garmanni 933, 938–939
variabilis 155–161
variabilis marmoratus 156, 158
variabilis olloporus 156, 158
variabilis smithi 156, 158
variabilis teapensis 156
variabilis variabilis 156–159
virgatus 338–839
Scirpus cyperinus 608
Scomber scomber 375
Scomberesox 738
Scombrosphyraena 780
Semaprochilodus 618
Semotilus 112, 1076
atromaculatus 538, 625, 822
margarita 822
Seriphus politus 691–696
Serrasalmus 618
Setipinna 53
Sibon 964 *sanniola* 963–969
Siebenrockiella crassicollis 920, 923
Simoselaps 200
Siphonognathus 95
Siphonops syntremus 861–862
Siren 627 *intermedia* 630
Sitana ponticeriana 477, 485
Skiffia 498, 505, 508–509
bilineata 499–500, 505–506, 508
(*Neotoca*) *bilineata* 505
francesae 499–500, 504–505, 508
(*Ollentodon*) *francesae* 505
lermae 499–500, 505–506, 508–509
(*Skiffia*) *lermae* 505
multipunctata 499–500, 505–506, 508
(*Ollentodon*) *multipunctata* 505
variegata 508
Smilisca 644
baudinii 641, 643–644, 646
phaeota 641, 643–644
puma 641, 643–644, 646
Solenopsis invicta 208
Sonora episcopa 965, 967
Sparisoma viride 830
Spartina
alterniflora 715, 717, 719, 722, 724
parens 715, 719, 724
Spermophilus
columbianus 478
richardsoni 478
Sphaenorhynchus 641 *carneus* 641, 643
Sphaerodactylus 466
macrolepis 466
parkeri 466
rosaurae 466
Sphenomorphus spp. 201
Sphyraenops 780
Staurotypus 925

salvini 924–925
triporcatus 923
Stenodactylus 467–468, 473, 477
Stephanolepis hispidus 331–338
Stereolepis 781 *gigas* 778
Sternotherus 925–926
carinatus 924
minor 924, 926
odoratus 264–268, 924
Stizostedion
canadense 249
lucioperca 249
vitreum 246–250
Stolephorus 48, 53
buccaneeri 52–53
curtus 51
exiguus 49, 51
indicus 48, 50, 53
ischanus 49, 51
lucida 51
miarchus 49
mundeolus 48, 50–52
naso 51
scofieldi 51
spp. 53
starksi 51
Strophiurichthys 902, 912–917
inermis 916
robustus 904, 910, 916
Strophurus 168
Sus 479
Suttonia 784 *lineata* 778
Syllis spp. 334
Sylvilagus bachmani 512
Syngnathus spp. 830
Synurella chamberlaini 631
Syringodium filiforme 1014, 1019, 1022

Tapatia 497
Tarentola 467–468, 473
Telmatobius 30, 34–36
Telmatobufo 35
Tenebrio 701 sp. 836
Teratolepis 467, 473
Teratoscincus 467, 473
Terrapene
carolina 922
carolina triunguis 922
coahuila 922
ornata 215, 218–219, 921
Testudo
gigantea 215
graeca 923
hermanni 923
Thalassia testidinum 333, 336
Thalassoma 826, 828
bifasciatum 255, 830
cupido 826, 828
pavo 826
"*Thalassoma*" 826, 828 sp. 827
Thamnophis 511–512, 1093
couchi 414
crytopsis 586
elegans 511–513, 1092–1096
elegans vagrans 262–264
melanogaster 530–534
melanogaster canescens 530, 532–533
melanogaster chihuahuaensis 533
melanogaster linearis 532
sauritus 1097
sirtalis 211, 965, 967, 1036–1041, 1095, 1097
Thecadactylus 467, 473
Thomomys umbrinus 76
Thorius 1051–1073
dubitus 1052–1053, 1055, 1057, 1063–1067, 1070
macdougalli 1052–1053, 1055, 1062, 1064–1070
maxillabrochus 1051, 1053, 1057, 1067–1069
minutissimus 1053, 1055–1056, 1065–1066, 1069
narisovalis 1052–1053, 1055, 1063–1066
pennatulus 1052–1053, 1062, 1067, 1070
pennatulus narismagnus 1052, 1055, 1065–1066
pennatulus pennatulus 1051, 1055, 1065
pulmonaris 1052–1053, 1056, 1063–1066
schmidti 1051, 1053, 1055, 1057, 1063–1066, 1068–1070
sp. A 1055, 1064–1065, 1068, 1070
sp. B 1055, 1064–1065, 1068, 1070
sp. C 1055, 1064–1065, 1068–1070
sp. D 1055, 1064–1065, 1068–1070
sp. E 1055, 1064–1065, 1068, 1070
sp. F 1056, 1065, 1068–1069
sp. G 1056, 1064–1065, 1068
troglodytes 1052–1053, 1055, 1057, 1062, 1064–1067, 1070
Thrissina 53 *encrasicholoides* 48, 50
Thryssa 53
Tilapia 753
aurea 753
macrocephala 961
Tomistoma 89, 92–94, 844
machikanense 89–95
schlegeli 89, 92–94
taiwanicus 94
Toxicodendron diversiloba 511
Toyotamaphimeia 89–95 *machikanensis* 89–95
Trachelochismus 865
melobesia 863
pinnulatus 863
sp. 863
Trachydactylus 473
Tribolium 815
Trichodon trichodon 549–551
Trichosurus vulpecula 156
Trifolium sp. 992
Triodon 905, 914
Trionyx 926
cartilagineus 925
ferox 925, 1087
leithii 926
muticus 925

sinensis 925
spiniferus 925
Tripterygion tripteronotus 833
Trisotropis dermopterus 778
"Trisotropis" dermopterus 784
Triturus 662–672
alpestris 669–670
alpestris apuanus 663, 667–669
alpestris group 669
boscai 662–672
boscai × *Triturus alpestris apuanus* 667
boscai × *Triturus vulgaris meridionalis* 667
carnifex 670
cristatus 668–670
cristatus carnifex 669
cristatus cristatus 669
cristatus dobrogicus 669
cristatus group 669–670
cristatus karelinii 669
dobrogicus 670
helveticus 669–670, 983
italicus 667, 669–670
italicus × *Triturus boscai* 667
karelinii 670
marmoratus 669
montandoni 669–670
vittatus 662–672
vittatus × *Triturus vulgaris* 668
vittatus ophryticus 662–663, 667–669
vittatus ophryticus × *Triturus alpestris apuanus* 663, 669
vittatus ophryticus × *Triturus boscai* 669
vittatus ophryticus × *Triturus cristatus carnifex* 669
vittatus ophryticus × *Triturus vittatus meridionalis* 669
vittatus ophryticus × *Triturus vittatus vulgaris* 669
vittatus vittatus 663
vulgaris 668–670
vulgaris group 669–670
vulgaris meridionalis 667, 669
vulgaris meridionalis × *Triturus boscai* 667
vulgaris vulgaris 669
Tropidodipsas 964
Tropidurus 131, 134, 139
albemarlensis 139
hispidis 139
torquatus 131–141
Tropiocolotes 467–469, 472–473, 477
depressus 468, 472
helenae 468–472
heteropholis 468, 472
latifi 468–472
persicus 468, 472
scortecci 468, 472
steudneri 468, 472
tripolitanus 468, 472
Tupinambis
nigropunctatus 477–478
teguixin 477
Typha latifolia 608
Typhlonectes natans 862
Typhlops brongersmianus 799
Ulmus rubra 340
Umbra limi 417
Underwoodisaurus 470
Urocotyledon 472
inexpectata 471–472
palmata 471–472
Uromastyx spinipes 477
Uroplatus 467, 473
Urosaurus 597
graciosus 139
ornatus 585–598, 741
Uta stansburiana 433, 586–587, 744

Vallisneria sp. 544
Varanus
bengalensis 477
griseus 477
komodoensis 477
prasinus 477
salvator 477
timorensis 477
Variola 784 *louti* 778
Vermicella 200

Wasmannia auropunctata 322
Wetmorella 875–883
albofasciata 875, 878–879, 881–882
nigropinnata 875–876, 878–879, 881–882
ocellata 875, 879–881
philippina 875–876, 878–879, 881
philippina bifasciata 875, 879–882
philippina philippina 875
phillipina 879
triocellata 875, 879, 881

Xanthosoma 871–872
Xantusia vigilis 158, 586
Xenobalistes tumidipectoris 906
Xenochrophis vittatus 967
Xenoophorus 498, 503, 505, 507, 509
captivus 499–500, 506–507
erro 507
Xenopus 420, 427 *laevis* 411–420, 536
Xenotaenia 498, 507 *resolanae* 499–500, 506–507
Xenotoca 498, 503, 507, 509
eiseni 499–500, 506–507
melanosoma 499–500, 506–507
variata 499–500, 506–507
Xiphophorus 171, 175–178, 180, 756–757, 766–768, 975, 978
clemenciae 766
cortezi 756, 766
helleri 756, 766–767, 975, 977
maculatus 170–181, 756, 758, 764, 766–768, 770–774
maculatus × *Xiphophorus helleri* 179
milleri 756, 766
montezumae 755–769
nigrensis 177, 756, 766, 768

pygmaeus 177, 756, 766
variatus 766, 975, 977
xiphidium 756, 766

Youngina 485

Zonotrichia leucophrys nuttalli 511–513
Zoogoneticus 498, 507 *quitzeoensis* 499–500, 506–507

SUBJECT INDEX

COPEIA, 1983, NOS. 1–4

ACCLIMATION, *Anolis carolinensis* (& evaporative water loss) 701–704.

ACTIVITY, *Plethodon websteri* (seasonal activity) 339–346; *Rana catesbeiana* (oxygen consumption during forced activity) 407–410; *Chelonia mydas* (seasonal & diel activity in a Florida lagoon) 1013–1023; *Podarcis taurica ionica* (diurnal & seasonal) 991–1001.

AGE, *Xiphophorus maculatus* (age/weight relationship to sexual maturation) 770–774; *Sceloporus o. occidentalis* (age, size, maturation relationships to home range) 933–940.

BEHAVIOR, *Rana clamitans* (agonistic behavior to calls) 141–155; *Diplodactylus spinigerus* (ejection of caudal gland secretion) 161–169; *Hoplocephalus* (*bitorquatus, bungaroides, stephensi*) (arboreality) 198–205; *Cnemidophorus sexlineatus* (hibernation) 206–214; *Nomeus gronovii* (commensal relationship with *Physalia physalis*) 250–252; *Scarus vetula* (spawning behavior) 253–256; *Xenopus laevis* (influence of social & predatory stimuli on air-breathing behavior) 411–420; *Phyllomedusa vaillanti* (social aggregations, feeding) 420–428; *Caretta caretta* (tidal cycle & emergence from sea) 516–519; *Crotalus* (*durissus, viridis viridis*) (chemosensory searching) 519–522; *Apeltes quadracus, Gasterosteus aculeatus* (interspecific aggression) 541–544; *Cyrtocara* (*annectens, moorii*), *Lethrinops praeorbitalis* (feeding & aggression) 544–547; resin-coated fish models for behavioral studies 547–549; *Trichodon trichodon, Oncorhynchus gorbuscha* (mixed schools of juveniles in near-shore areas) 549–551; *Gambusia* (*puncticulata yucatana, sexradiata*) (behavioral separation of coexisting spp.) 598–607; *Gramma loreto* (territoriality) 829–832; *Forsterygion varium* (homing) 832–834; *Crotaphytus collaris* (effects of movement & morphol. on predation) 835–837; *Gephyrosaurus bridensis* (inferred behavior from mandibular fractures) 845–847; *Ambystoma jeffersonianum, Bolitoglossa subpalmata, Eurycea bislineata, Notophthalmus viridescens, Plethodon jordani* (antipredator behavior) 1036–1041; *Lepomis macrochirus* (mobbing of snapping turtle as antipredator behavior) 1086–1088; *Desmognathus ochrophaeus* (aggression towards conspecifics) 1098–1101.

BLOOD, *Pseudemys scripta elegans, Chrysemys picta belli* (collecting by cardiac puncture) 522–523.

BREATHING, *Xenopus laevis* (influence of social & predatory stimuli on air-breathing) 411–420.

BREEDING, *Platynotus semitaeniatus, Tropidurus torquatus* (seasonality factors) 131–141; *Ambystoma t. tigrinum* (breeding migration) 608–616; *Fundulus grandis* (season) 711–718.

CALLS, *Rana clamitans* (size-related variation) 141–155; *Ranidella riparia* 534–537; neotropical hylid frogs (acoustic resource partitioning) 639–649; *Eleutherodactylus* frogs (acoustic community characteristics) 941–953; *Eleutherodactylus* (*coqui, portoricensis*) (acoustic communication) 1089–1090.

CANNIBALISM, *Ambystoma tigrinum mavortium* (morphs) 1006–1012.

CAUDAL GLANDS, *Diplodactylus spinigerus* (secretions, histol., morphol., evol. of) 161–169.

CHEMORECEPTION, *Crotalus* (*durissus, viridis viridis*) (odors) 519–522; *Chrosomus neogaeus, Notropis* (*venustus, whipplei*) (seasonal loss of alarm substance) 822–826.

CHROMOSOMES, *Insuetophrynus acarpicus* (karyotypes) 30–37; *Hyla chrysoscelis* (chromosome polymorphism) 273–275; *Salmo gairdneri* (chromosomal evol.) 650–662; *Triturus* (*boscai, vittatus*) (karyology) 662–672; *Xiphophorus montezumae* (sex-determining mechanism) 755–769; cryptodiran turtles (karyology) 918–932; *Alsodes* (*barrioi, gargola, monticola, nodosus, tumultuosus, vanzolinii, verrucosus*) (karyological relationships), 1104–1107.

COLORATION, *Sceloporus variabilis* (unilateral testicular pigmentation) 155–161; *Xiphophorus maculatus* (genetic basis of mutant color varieties) 170–181; *Thamnophis* (*canescens, melanogaster*) (color & pattern variation) 530–534; *Ptychadena mascareniensis, Rana* (*clamitans, pipiens, sylvatica*) (dark-pigment variants, color changes, descriptive guidelines) 979–990.

COMPETITION, *Notophthalmus viridescens dorsalis, Ambystoma tigrinum* (competitive interactions) 628–639.

CYTOLOGY, *Ctenopharyngodon idella* (gonadal differentiation) 749–755.

DEVELOPMENT, *Cyprinus carpio, Carassius auratus* (larval development) 116–121; *Stizostedion vitreum, Catostomus commersoni* (form & function during embryonic development) 246–250; *Peprilus* (*burti, paru, triacanthus*) (larval development) 397–406; *Rana catesbeiana* (metamorphosis and oxygen consumption) 407–410; *Gobiesox maeandricus* (larval development) 551–554; *Cyprinus carpio* (of clonal state in successive gynogenetic generations) 745–749; *Poecilia formosa* (embryonic development, superfetation) 969–974.

DISTRIBUTION, *Lophiomus setigerus, Lophius* (*americanus, budegassa, gastrophysus, litulon, piscatorius, upsicephalus, vaillanti, vomerinus*) (geogr. & bathymetric distr.) 11–30; *Fundulus zebrinus* 55–67; *Oryzias latipes* (genetic variation & geogr. popns.) 311–318; *Monacanthus ciliatus, Stephanolepis hispidus* (temporal & spatial distr. in Apalachee Bay,

Fla.; pollution effects) 331–338; *Phyllomedusa vaillanti* (spatial, diel patterns) 420–428; *Anolis limifrons* (dispersal patterns, home range) 429–434; *Bagarius* (*bagarius, suchus* sp. nov., *yarrelli*) (geogr.) 435–445; *Noturus* (*baileyi, flavipinnis*) (discovery in Citico Cr., Little Tennessee R. system) 558–560; *Poecilia* (*Mollienesia*) spp. (spp. ranges) 817–822; *Sceloporus o. occidentalis* (home range) 933–940; *Eleutherodactylus* frogs (spatial patterns of acoustic communities) 941–953; *Etheostoma* (*euzonum erizonum, e. euzonum*) 1041–1050.

DUVERNOY'S GLAND, *Thamnophis elegans vagrans* (function of) 262–264.

ECOLOGY, *Insuetophrynus acarpicus* 30–37; *Gambusia affinis* (allele frequencies, environ. variables) 182–193; *Hoplocephalus* (*bitorquatus, bungaroides, stephensi*) 198–205; *Anolis opalinus* (resource partitioning) 319–331; *Plethodon websteri* 339–346; *Gambusia* (*puncticulata yucatana, sexradiata*) 598–607; neotropical hylid frogs (acoustic resource partitioning) 639–649; *Gastrotheca* (*bufona, cornuta, dendronastes*) (ecol. of trans-Andean marsupial frogs) 868–874; *Podarcis taurica ionica* (habitat ecol.) 991–1001; *Thorius* (*dubitus, macdougalli, minutissimus, narisovalis, pennatulus pennatulus, pennatulus narismagnus, pulmonaris, schmidti, troglodytes,* spp. undescribed) (ecol. relationships) 1051–1073.

EGGS, *Chelydra* (*picta, serpentina*), *Chrysemys picta, Emydoidea blandingi, Graptemys* (*ouachitensis, geographica*), *Pseudemys scripta, Sternotherus odoratus* (egg components) 264–268; *Chelydra serpentina, Geochelone elephantopus, Lepidochelys kempi,* and fossil chelonians (fossil & contemporary eggshell structure) 382–397; *Hyla* (*cinerea, gratiosa, regilla*) (multiple clutches) 513–516; *Urosaurus ornatus* (clutch & egg size) 585–598; *Fundulus heteroclitus* (distr. in relation to tidal height) 719–725; *Fundulus heteroclitus* (morphol. of egg with geogr.) 726–740; *Rana sylvatica* (natural survival of eggs & larvae) 1096–1098.

ELECTROPHORESIS, *Crotaphytus collaris baileyi* × *C. insularis bicinctores* (hybridization in Coconino Co., Ariz.) 1–11; *Insuetophrynus acarpicus* (eyelens) 30–37; *Catostomus discolobus yarrowi* (tissue & serum) 37–48; *Ambystoma* (*rosaceum, tigrinum*) 67–78; *Ranidella* (*riparia, signifera*) (allele frequencies) 275–279; *Oryzias latipes* 311–318; *Anolis carolinensis* 523–529; *Lucania* (*goodei, interioris, parva*) (electrophoretic variation) 564–570; *Micropterus* (*dolomieui, treculi*) (allozyme loci) 672–679; *Dicamptodon* (*aterrimus, copei, ensatus*) 679–691; *Pseudoeurycea* (*altamontana, leprosa, longicauda* sp. nov., *robertsi*) (compar. of allozymes) 884–894; *Thorius* (*dubitus, macdougalli, minutissimus, narisovalis, pennatulus narismagnus, pennatulus pennatulus, pulmonaris, schmidti, troglodytes,* spp. undescribed) (analyses of spp. & subspp.) 1051–1073.

EMBRYOLOGY, *Stizostedion vitreum, Catostomus commersoni* (compar. embryonic ecomorphol.) 246–250; *Isurus paucus, Alopias superciliosus* (embryos, oophagy) 375–382; *Poecilia formosa* (superfetation, embryonic development) 969–974.

EVOLUTION, *Diplodactylus spinigerus* (evol. of caudal glands) 161–169; *Micropterus salmoides* (rapid evol. in post-thermal environ.) 192–197; *Salmo gairdneri* (chromosomal evol.) 650–662; *Dicamptodon* (*aterrimus, copei, ensatus*) 679–691; cryptodiran turtles (karyotypic evol.) 918–932.

FECUNDITY, *Ichthyomyzon* (*castaneus, gagei*) 367–374.

FEEDING, *Chelonia mydas* (popns. with divergent nesting locations but shared feeding ranges in Brazil) 121–125; *Alopias superciliosus, Isurus paucus* (embryonic oophagy) 375–382; *Phyllomedusa vaillanti* (diel patterns of) 420–428; *Sceloporus virgatus* (assimilation efficiency) 838–839; *Chelonia mydas* (feeding ecol.) 1013–1023.

FOOD, *Hoplocephalus* (*bitorquatus, bungaroides, stephensi*) (food habits) 198–205; *Hyla chrysoscelis, Rana clamitans* (food quality & quantity & early larval growth) 238–242; *Alopias superciliosus, Isurus paucus* (embryonic oophagy) 375–382; *Anolis opalinus* (diet, food-resource partitioning) 319–331; *Monacanthus ciliatus, Stephanolepis hispidus* (food habits, effects of pulp-mill pollution on feeding habits) 331–338; *Phyllomedusa vaillanti* (habits) 420–428; *Hemigaleus microstomus* (diet) 487–497; *Thamnophis elegans* (diet) 511–513; *Gambusia* (*puncticulata yucatana, sexradiata*) (compar. of diets) 598–607; stat. analysis on food & growth relationships 1090–1092.

FUNCTION, *Agkistrodon piscivorus, Constrictor constrictor, Crotalus viridis, Naja naja, Pituophis melanoleucus* (function of rectus capitus muscle in biting) 221–225; *Thamnophis elegans vagrans* (function of Duvernoy's gland) 262–264; lacertilian hind limb (& locomotion) 476–487.

GENETICS, *Xiphophorus maculatus* (genetic basis of mutant color varieties) 170–181; *Gambusia affinis* (biochemical genetics, allelic variation) 182–193; *Micropterus salmoides* (gene frequency & thermal environ.) 193–197; *Ranidella* (*riparia, signifera*) (genetic divergence between) 275–279; *Poeciliopsis lucida* (genetic heterozygosity & meristic variation) 287–299; Goodeidae (karyology) 497–510; *Anolis carolinensis* 523–529; *Micropterus* (*dolomieui, treculi*) (genetic introgression) 672–679; *Genyonemus lineatus, Paralabrax clathratus, Seriphus politus* (popn. genetic structure) 691–696; *Scaphirhynchus* (*albus, platorynchus*) (genetic similarity) 696–700; *Cyprinus carpio* (development of clonal state in successive gynogenetic generations) 745–749; *Xiphophorus montezumae* (sex-determining mechanism) 755–769; *Xiphophorus maculatus* (of sexual

maturation) 770–774; *Ptychadena mascareniensis, Rana (clamitans, pipiens, sylvatica)* (inheritance of color patterns) 979–990.

GEOGRAPHIC LOCALITIES

Africa, *Cyrtocara (annectens, moorii), Lethrinops praeorbitalis)* (Lake Malawi) 544–547.

Alabama, *Campostoma (oligolepis, pauciradii* sp. nov.) 101–116; *Cnemidophorus sexlineatus* 206–214; *Ichthyomyzon gagei* 367–374; *Anolis carolinensis* 523–529; *Fundulus grandis* 711–718.

Alaska, *Oncorhynchus gorbuscha, Trichodon trichodon* 549–551; *Salmo gairdneri* 650–662.

Argentina, *Minyichthys inusitatus* sp. nov. 774–776; *Liophis miliaris* 791–802.

Arkansas, *Opheodrys aestivus* 741–745; *Etheostoma (euzonum erizonum, e. euzonum)* 1042–1050.

Arizona, *Crotaphytus collaris baileyi* × *C. insularis bicinctores, Catostomus (d. discolobus, discolobus yarrowi)* 37–48; *Urosaurus ornatus* 585–598; *Sceloporus virgatus* 838–839.

Australia, *Isurus oxyrinchus* (New South Wales) 126–130; *Hoplocephalus (bitorquatus, bungaroides, stephensi)* 198–205; *Eretmochelys imbricata* (Torres Strait) 271–273; *Ranidella (riparia, signifera)* (W. Austr.) 275–279; *Oedura (gemmata* sp. nov., *marmorata)* 445–449; *Hemigaleus microstoma* 487–497; *Ranidella riparia* (W. Austr.) 534–537; *Gambusia affinis* (W. Austr.) 953–963.

Bahamas, *Gramma loreto* 829–832.

Bangladesh, *Bagarius bagarius* 435–442.

Belize, *Gambusia (puncticulata yucatana, sexradiata)* 598–607.

Bolivia, *Liophis miliaris* 791–802.

Borneo, *Bagarius yarrelli* 435–445.

Brazil, *Chelonia mydas* 121–125; *Platynotus semitaeniatus, Tropidurus torquatus* 131–141; *Cnemidophorus ocellifer* (ne. Brazil) 359–366; *Phyllomedusa vaillanti* 420–428; neotropical hylid frogs 639–649; *Liophis miliaris* 791–802.

Burma, *Bagarius yarrelli* 435–442.

California, *Anniella (nigra argentea, nigra nigra)* 79–89; *Dipsosaurus dorsalis* 268–271; *Xenopus laevis* 411–420; *Thamnophis elegans* 511–513; *Hyla regilla* 513–516; *Salmo gairdneri* 650–662; *Dicamptodon ensatus* 679–691; *Genyonemus lineatus, Paralabrax clathratus, Seriphus politus* 691–696; *Sceloporus o. occidentalis* 933–940; *Thamnophis elegans* 1092–1096.

Canada, *Salmo gairdneri* (B.C.) 650–662; *Coryphopterus nicholsi* (B.C.) 809–812; *Chrosomus neogaeus* (Saskatchewan) 822–826.

Canary Islands, *Coryphaenoides thelestomus* 1083–1086.

Caribbean, *Minyichthys inusitatus* sp. nov. 774–776.

Chile, *Insuetophrynus acarpicus* 30–37; *Liolaemus fabiani* 788–790; *Liophis miliaris* 791–802; *Alsodes (vanzolinii, verrucosus)* 1104–1107.

Colombia, *Liophis miliaris* 791–802; *Amphignathodon guentheri, Gastrotheca (angustifrons, bufona, cornuta, dendronastes)* 868–874.

Connecticut, *Fundulus heteroclitus* 726–740.

Costa Rica, neotropical hylid frogs 639–649; *Bolitoglossa subpalmata* 1036–1041.

Delaware, *Fundulus heteroclitus* 719–725.

Ecuador, neotropical hylid frogs 639–649; *Liophis miliaris* 791–802; *Amphignathodon guentheri, Gastrotheca (angustifrons, cornuta, dendronastes)* 868–874.

Fiji, *Paracaesio stonei* sp. nov. 450–453.

Florida, *Gambusia affinis, Poecilia (latipinna, reticulata)* 243–245; *Nomeus gronovii* 250–252; *Monacanthus ciliatus, Stephanolepis hispidus* 331–338; *Alopias superciliosus, Isurus paucus* 375–382; *Anolis carolinensis* 523–529; *Lucania (goodei, parva)* 564–570; *Fundulus heteroclitus* 726–740; *Chelonia mydas* 1013–1023.

French Guiana, *Liophis miliaris* 791–802.

Georgia, *Campostoma (anomalum michauxi, oligolepis, pauciradii)* 101–116; *Gambusia affinis* 182–193; *Cnemidophorus sexlineatus* 206–214; *Hyla (cinerea, gratiosa, regilla)* 513–516; *Caretta caretta* 516–519.

Grand Cayman Is., BWI, *Gambusia xanthosoma* sp. nov. 457–464.

Great Britain, *Gephyrosaurus bridensis* 845–847.

Greece, *Podarcis taurica ionica* 991–1001.

Guatemala, *Minascaecilia sartoria* gen. nov., sp. nov. 857–863.

Gulf of Mexico, *Gonostoma elongatum* 554–557.

Guyana, *Liophis miliaris* 791–802.

Idaho, *Salmo gairdneri* 650–662; *Dicamptodon aterrimus* 679–691.

Illinois, *Acris crepitans* 300–311; *Scaphirhynchus platorynchus* 696–700; *Notropis whipplei* 822–826.

India, *Bagarius yarrelli* 435–445.

Indiana, *Ambystoma texanum* 624–628.

Iowa, *Acris crepitans* 300–311.

Jamaica, *Anolis opalinus* 319–331.

Japan, *Tomistoma machikanense* 89–95; *Oryzias latipes* 311–318.

Java, *Bagarius yarrelli* 435–445.

Kentucky, *Ambystoma texanum* 624–628.

Laos, *Bagarius yarrelli* 435–445.

Louisiana, *Anolis carolinensis* 523–529; *Notropis venustus* 822–826.

Maine, *Fundulus heteroclitus* 726–740.

Maryland, *Cyprinus carpio, Carassius auratus* 116–121; *Fundulus heteroclitus* 726–740; *Ambystoma (jeffersonianum, maculatum)* 1001–1005.

Massachusetts, *Fundulus heteroclitus* 726–740.

Mexico, *Ambystoma (rosaceum, triginum)* (nw, Sierra Madre Occidental) 67–78; *Anniella (pulchra, nigra argentea)* (Baja Calif.) 79–89; Goodeidae 497–510; *Thamnophis (canescens, melanogaster)* (Michoacan) 530–534; *Lucania parva* (Yucatan) 564–570; *Xiphophorus montezumae* 755–769; *Lutjanus guttatus* (Sonora) 813–816; *Pseudoeurycea (altamontana, leprosa, longicauda* sp. nov., *robertsi)* 884–894; *Sibon sanniola* (Yucatan Pen.) 963–969; *Poecilia formosa* (Tamaulipas) 969–974; *Thorius* spp. (Oaxaca, Veracruz, Puebla) 1051–1073.

Michigan, *Chelydra serpentina, Emydoidea blandingi* 264–268; *Ichthyomyzon castaneus* 367–374.

Mississippi, *Campostoma anomalum pullum* 101–116; *Ioa vigil, Percina shumardi* 1082–1083.
Missouri, *Cryptobranchus alleganiensis bishopi* 225–231; *Scaphirhynchus* (*albus, platorynchus*) 696–700; *Etheostoma* (*euzonum erizonum, e. euzonum*) 1042–1050.
Montana, *Scaphirhynchus platorynchus* 696–700.
Nebraska, *Scaphirhynchus platorynchus* 696–700.
Nepal, *Rana* (*Paa*) (*annandalii, blandfordii, polunini, rara* sp. nov., *rostandi*) 895–901.
New Guinea, *Litoria umbonata* sp. nov. 803–808.
New Hampshire, *Osmerus mordax* 812–813.
New Mexico, *Catostomus* (*discolobus yarrowi, plebius*) 37–48.
New Jersey, *Fundulus heteroclitus* 726–740; *Rana sylvatica* 1096–1098.
New Mexico, crocodilian fossil 842–846; *Boleosoma phlox, Etheostoma spectabile pulchellum* 1082–1083.
New York, *Apeltes quadracus, Gasterosteus aculeatus* 541–544; *Lepomis macrochirus* 1086–1088; *Ambystoma jeffersonianum, Eurycea bislineata, Notophthalmus viridescens, Thamnophis sirtalis* 1036–1041.
New Zealand, *Odax* (*cyanoallix* sp. nov., *pullus*) 95–101; *Forsterygion varium* 832–834; *Modicus* gen. nov. (*minimus* sp. nov., *tangaroa* sp. nov.) 863–868.
North Atlantic, *Peprilus* (*burti, paru, triacanthus*) 397–406; *Coryphaenoides* (*alateralis, thelestomus*) (off Fla. & NJ) 1083–1086.
North Carolina, *Campostoma* (*anomalum michauxi, oligolepis*) 101–116; *Hyla gratiosa* 232–237; *Hyla chrysoscelis, Rana clamitans* 238–242; *Notophthalmus viridescens dorsalis, Ambystoma tigrinum* 628–639; *Fundulus heteroclitus* 726–740; *Plethodon jordani* 1036–1041; *Desmognathus ochrophaeus* 1098–1101.
Oregon, *Salmo* sp. (fossil) 563–564; *Salmo gairdneri* 650–662; *Dicamptodon* (*copei, ensatus*) 679–691.
Ohio, *Ambystoma texanum* 624–628.
Oklahoma, *Crotaphytus collaris* 835–837.
Pacific (eastern), *Anchoa* (*argentivittata, duodecim, mundeola, panamensis*), *Anchoviella* (*miarcha, scitula*) 48–54.
Panama, *Anolis limifrons* 429–434; *Pseudemys scripta* 529–530; *Iguana iguana* 705–711.
Paraguay, *Loricaria simillima, Loricariichthys* (*labialis, platymetopon*) 257–259; *Liophis miliaris* 791–802.
Pennsylvania, *Carassius auratus, Cyprinus carpio,* 116–121; *Notropis atherinoides* × *N. volucellus* 1077–1082; *Desmognathus f. fuscus* 1101–1104.
Puerto Rico, *Scarus vetula* 253–256; *Eleutherodactylus* frogs 941–953; *Eleutherodactylus* (*coqui, portoricensis*) 1089–1090.
South Carolina, *Campostoma anomalum michauxi* 101–106; *Gambusia affinis,* 182–193; *Micropterus salmoides* 193–197; *Pseudemys scripta* 264–268; *Plethodon websteri* 339–346; *Ambystoma t. tigrinum* 608–616; *Fundulus heteroclitus* 726–740.
South and Southeast Asia, *Bagarius* (*bagarius, suchus* sp. nov., *yarrelli*) 435–445.
Sumatra, *Bagarius yarrelli* 435–445.
Surinam, *Liophis miliaris* 791–802; *Chelonia mydas* × *Eretmochelys imbricata* 839–842.
Tennessee, *Campostoma oligolepis* 101–116; *Anolis carolinensis* 523–529; *Moxostoma atripinne* 538–541; *Noturus* (*baileyi, flavipinnis*) 558–560; *Hybopsis monacha* × *Notropis galacturus* 1074–1077.
Texas, *Anolis carolinensis* 523–529; *Lucania parva* 564–570; *Micropterus* (*dolomieui, treculi*) 672–679; *Poecilia formosa* 969–974; *Boleosoma phlox, Etheostoma spectabile pulchellum* 1082–1083.
Thailand, *Bagarius* (*bagarius, suchus* sp. nov., *yarrelli*) 435–445.
Uruguay, *Liophis miliaris* 791–802.
Venezuela, *Prochilodus mariae* 617–623.
Vietnam, *Bagarius yarrelli* 435–445.
Virginia, *Fundulus heteroclitus* 726–740.
Washington, *Gobiesox maeandricus* 551–554; *Salmo gairdneri* 650–662; *Dicamptodon* (*copei, ensatus*) 679–691.
Wisconsin, *Chelydra picta, Graptemys ouachitensis* 264–268; *Scaphirhynchus platorynchus* 696–700.
GROWTH, *Cryptobranchus alleganiensis bishopi* (growth rates) 225–231; *Hyla gratiosa* (larval growth & survival) 232–237; *Hyla chrysoscelis, Rana clamitans* (food influence on early larval growth) 238–242; *Gambusia affinis, Poecilia* (*latipinna, reticulata*) (salinity & size-hierarchy effect) 243–245; *Plethodon websteri* 339–346; *Gambusia affinis* (somatic & reprod. growth variation in experimental popns.) 953–963; stat. analysis on food & growth relationships 1090–1092; *Plethodon jordani* 1024–1035.

HABITAT, *Moxostoma atripinne* 538–541; *Gambusia* (*puncticulata yucatana, sexradiata*) (separation of coexisting spp.) 598–607; *Gramma loreto* (distr.) 829–832.
HERMAPHRODITISM, *Hemanthias peruanus* (protogynous) 252–253; *Gonostoma elongatum* (protandry) 554–557; *Coryphopterus nicholsi* (protogynous) 809–812; *Osmerus mordax* 812–813.
HIBERNATION, *Cnemidophorus sexlineatus* 206–214.
HISTOLOGY, *Diplodactylus spinigerus* (of caudal gland) 161–169.
HYBRIDIZATION, *Crotophytus collaris baileyi* × *C. insularis bicinctores* (Little Colorado R., Coconino Co., Ariz.) 1–11; *Campostoma pauciradii* × *C. oligolepis* 101–116; *Triturus* (*boscai, vittatus*) (hybridization experiments with *Triturus* spp.) 662–672; (*Micropterus dolomieui* × *M. treculi*) 672–679; *Chelonia mydas* × *Eretmochelys imbricata* 839–842; *Hybopsis monacha* × *Notropis galacturus* 1074–1077; *Notropis atherinoides* × *N. volucellus* 1077–1082.
LARVAE, *Cyprinus carpio, Carassius auratus* 116–121; *Peprilus* (*burti, paru, triacanthus*) (larval development) 397–406; *Gobiesox maeandricus* (larval development) 551–554; *Ambystoma texanum* (predation by fish) 624–628; "*Thalassoma*" larvae (occurrence in E. Pacific Barrier) 826–829; *Ambystoma* (*jeffersonianum, maculatum*) (temp. pref-

erences) 1001–1005; *Rana sylvatica* (natural survival of tadpoles) 1096–1098.

LIFE HISTORY, *Insuetophrynus acarpicus* 30–37; *Acris crepitans* 300–311; *Plethodon websteri* 339–346; *Moxostoma atripinne* (food, reprod., growth) 538–541; *Urosaurus ornatus* (popn. biol.) 585–598; *Gambusia affinis* (variation among experimental popns.) 953–963; *Podarcis taurica ionica* 991–1001.

LOCOMOTION, lacertilian hind limb & locomotion 476–487; *Thamnophis elegans* (food ingestion & locomotory performance) 1092–1096.

METABOLISM, *Chrysemys picta* (activity metabolism) 214–221.

MIGRATION, *Ambystoma t. tigrinum* (breeding migration) 608–616, *Prochilodus mariae* (usage of energy reserves during migration) 617–623

MORPHOLOGY, *Lophiomus, Lophius* (esca) 11–30; *Toyotamaphimeia machikanensis* (skull) 89–95; *Diplodactylus spinigerus* (caudal gland) 161–169; *Agkistrodon piscivorus, Constrictor constrictor, Crotalus viridis, Naja naja, Pituophis melanoleucus* (rectus capitus muscle) 221–225; *Chelydra serpentina, Geochelone elephantopus, Lepidochelys kempi,* fossil chelonians (eggshell morphol.) 382–397; *Cyrtodactylus* spp., *Microgecko* spp., *Nactus* gen. nov. (spp.), *Phyllodactylus* spp., *Pristurus* spp., *Tropiocolotes* spp., *Urocotyledon* gen. nov. (spp.) (branchial arch) 465–475; lacertilians (hind limb morphol.) 476–487; *Fundulus heteroclitus* (egg morphol. variation with geogr.) 726–740; *Pseudoeurycea (altamontana, leprosa, longicauda* sp. nov., *robertsi*) (morphometric analyses) 884–894.

MYOLOGY, Ostraciidae (aracanin genera) 902–917.

NESTING, *Chelonia mydas* (popns. with shared nesting ranges in Brazil) 121–125; *Pseudemys scripta* (island nesting & predation) 529–530; *Iguana iguana* (nest structure) 705–711; *Desmognathus ochrophaeus* (nest defense) 1098–1101.

NOMENCLATURE, *Fundulus zebrinus* (nomenclatural status) 55–67; *Toyotamaphimeia* gen. nov. 89–95.

ORGANS, *Thamnophis elegans vagrans* (Duvernoy's gland) 262–264; *Diplodactylus spinigerus* (caudal gland) 161–169.

OSTEOLOGY, *Cyrtodactylus* spp., *Microgecko* spp., *Nactus* gen. nov (spp.), *Phyllodactylus* spp., *Pristurus* spp., *Tropiocolotes* spp. *Urocotyledon* gen. nov. (spp.) (branchial arch) 465–475; Ostraciidae (aracanin genera) 902–917.

PALEONTOLOGY, *Toyotamaphimeia machikanense* (gen. nov.) (Pleistocene, central Japan) 89–95; *Salmo* sp. (Fossil Lake, Oregon) 563–564; crocodilian (early Paleocene endocranial cast) 842–846; *Gephyrosaurus bridensis* (mandibular fractures) 845–847.

PARASITISM, *Lutjanus guttatus* (tongue replacement by isopod) 813–816.

PHYSIOLOGY, *Chrysemys picta* (aerobic vs. anaerobic metabolism during activity) 214–221; *Xenopus laevis* (social & predatory stimuli influences on air-breathing) 411–420; *Anolis carolinensis* (acclimation & evaporative water loss) 701–704; *Sceloporus virgatus* (assimilation efficiency) 838–839.

POLLUTION, *Monacanthus ciliatus* (effects on distr. & feeding habits) 331–338.

POPULATION, *Catostomus (d. discolobus, discolobus yarrowi, plebeius)* (popn. in Ariz., N. Mex.) 37–48; *Fundulus zebrinus* (clinal variation) 55–67; *Ambystoma (rosaceum, tigrinum)* (nw. Mexico) 67–78; *Gambusia affinis* (biochem. variation in popns.) 182–193; *Dipsosaurus dorsalis* (popn. density in so. Calif.) 268–271; *Anolis carolinensis* (genetic variation & popn.) 523–529; *Urosaurus ornatus* (popn. biol.) 585–598; *Ambystoma t. tigrinum* (structure & dynamics during breeding) 608–616; *Dicamptodon (aterrimus, copei, ensatus)* (genetic variations in) 679–691; *Genyonemus lineatus, Paralabrax clathratus, Seriphus politus* (genetic structure) 691–696; *Podarcis taurica ionica* (density, sex ratio) 991–1001.

PREDATION, *Thamnophis elegans* (predation on white-crown sparrow) 511–513; *Pseudemys scripta* (island nesting & predation) 529–530; *Urosaurus ornatus* (predation) 585–598; *Ambystoma texanum* (by fish on larvae of) 624–628; *Ambystoma tigrinum, Notophthalmus viridescens dorsalis* (predatory interactions) 628–639; *Crotaphytus collaris* 835–837; *Ambystoma jeffersonianum, Bolitoglossa subpalmata, Eurycea bislineata, Notophthalmus viridescens, Plethodon jordani* (antipredator behavior) 1036–1041.

REPRODUCTION, *Insuetophrynus acarpicus* (oocytes) 30–37; *Isurus oxyrinchus* (in N.S.W., Austr.) 126–130; *Platynotus semitaeniatus, Tropidurus torquatus* (reprod. ecol.) 131–141; *Sceloporus variabilis* (reprod. activity & testicular pigmentation) 155–161; *Hoplocephalus (bitorquatus, bungaroides, stephensi)* (reprod. cycles, fecundities) 198–205; *Catostomus commersoni, Stizostedion vitreum* (reprod. guild classif.) 246–250; *Hemanthias peruanus* (protogynous hermaphrodite) 252–253; *Scarus vetula* (spawning behavior, mating systems) 253–256; *Loricaria simillima, Loricariichthys (labialis, platymetopon)* (reprod. ecol.) 257–259; *Gasterosteus aculeatus* (oocytes in testes) 259–261; *Eretmochelys imbricata* (reprod. migration) 271–273; *Plethodon websteri* 339–346; *Chrysemys picta* (seasonal gonadal cycle) 347–358; *Cnemidophorus ocellifer* 359–366; *Hemigaleus microstomus* (reprod. biol.) 487–497; *Caretta caretta* (tidal cycle & emergence from sea) 516–519; *Gonostoma elongatum* (protandric sex reversal) 554–557; *Eurypharynx pelecanoides* (re-

prod. strategy) 560–563; *Prochilodus mariae* (gonadal develop.) 617–623; *Fundulus grandis* (reprod. cycles & strategies) 711–718; *Fundulus heteroclitus* (spawning site use) 719–725; *Opheodrys aestivus* 741–745; *Ctenopharyngodon idella* (gonadal differentiation) 749–755; *Chrosomus neogaeus, Notropis (venustus, whipplei)* (spawning & seasonal loss of alarm substance) 822–826; *Gambusia affinis* (somatic & reprod. growth variation in experimental popns.) 953–963; *Sibon sanniola* (reprod. cycle in females) 963–969; *Podarcis taurica ionica* (age at maturity, egg size, clutch size, clutch per season) 991–1001; *Plethodon jordani* 1024–1035; *Desmognathus ochrophaeus* (brooding duration) 1098–1101;

RESPIRATION, *Xenopus laevis* (social & predatory stimuli influence on air breathing) 411–420; *Phyllomedusa vaillanti* (aerial; synchrony in) 420–428.

SEASONALITY, *Acris crepitans* (seasonal variation in color morph proportions) 300–311; *Anolis opalinus* (seasonal effect on diet) 319–331; *Monacanthus ciliatus, Stephanolepis hispidus* (seasonal overlap in distr.) 331–338; *Plethodon websteri* (seasonal activity & growth) 339–346; *Chrysemys picta* (seasonal gonadal cycle) 347–358; *Cnemidophorus ocellifer* (seasonality of abdominal fat bodies) 359–366; *Urosaurus ornatus* (annual cycles) 585–598; *Prochilodus mariae* (seasonal changes in caloric density of body compartments) 617–623; *Fundulus grandis* (of breeding cycles) 711–718; *Chrosomus neogaeus, Notropis (venustus, whipplei)* (seasonal loss of alarm substance) 822–826; *Sibon sanniola* (of reprod. cycle in females) 963–969; *Chelonia mydas* (of activity & movements in a Fla. lagoon) 1013–1023.

SENSORY CANAL, *Anchoa (argentivittata, duodecim, mundeola, panamensis), Anchoviella (miarcha, scitula)* 48–54.

SEX, *Ctenopharyngodon idella* (gonadal differentiation) 749–755; *Xiphophorus montezumae* (sex-determining mechanism) 745–769.

SEXUAL DIMORPHISM, *Cnemidophorus ocellifer* 359–366; *Gonostoma elongatum* (size dimorph.) 554–557; *Eurypharynx pelecanoides* (nasal rosette) 560–563; *Urosaurus ornatus* 585–598.

SIZE, *Hoplocephalus (bitorquatus, bungaroides, stephensi)* (size-freq., birth size, growth rates) 193–205; *Gambusia affinis, Poecilia (latipinna, reticulata)* (salinity & size-hierarchy effect) 243–245; *Plethodon websteri* 339–346; *Gonostoma elongatum* (dimorphism) 554–557; *Urosaurus ornatus* (size & age at maturity, sexual dimorph., age relationships, growth) 585–598; *Ambystoma t. tigrinum* (variation in body size) 608–616; *Sceloporus o. occidentalis* (age, size, maturation, relationsh ip to home range) 933–940; *Ambystoma tigrinum mavortium* (morphs, cranial shape & size) 1006–1012; *Desmognathus f. fuscus* (microgeogr. size variation) 1101–1104.

SPERMATOZOA, silver staining of, from living & preserved fishes, 974–978.

SURVIVORSHIP, *Urosaurus ornatus* 585–589; *Ambystoma texanum* (of larvae) 624–628; *Ambystoma tigrinum, Notophthalmus viridescens dorsalis* (predation & embryonic survival) 628–639; *Plethodon jordani* 1024–1035; *Ambystoma jeffersonianum, Bolitoglossa subpalmata, Eurycea bislineata, Notophthalmus viridescens, Plethodon jordani* (advantages of alternative antipredator strategies) 1036–1041; *Rana sylvatica* (of eggs & tadpoles) 1096–1098.

SYSTEMATICS, *Anchoa (argentivittatus, duodecim, mundeola, panamensis), Anchoviella (miarcha, scitula)* (status) 48–54; *Fundulus zebrinus* (status) 55–67; *Anniella (p. pulchra, n. nigra, n. argentea)* (nomenclature) 79–89; *Bagarius (bagarius, suchus* sp. nov., *yarrelli)* (revision of genus) 435–445; Oedura (*gemmata* sp. nov., *marmorata*) (new Australian species) 445–449; *Paracaesio stonei* sp. nov. (new Fijian species) 450–453; *Ctenoblepharis adituvelatus* sp. nov. (new Chilean sp.) 454–457; *Gambusia (beebei, eurystoma, luma, pseudopunctata, punctata, rhizophorae, sexradiata, xanthosoma* sp. nov.) (new sp. Grand Cayman Is.) 457–464; *Cyrtodactylus* spp., *Microgecko* spp., *Nactus* gen. nov. (spp.), *Phyllodactylus* spp., *Pristurus* spp., *Tropiocolotes* spp., *Urocotyledon* gen. nov. (spp.) (revision of gekkonids) 465–475; Goodeidae (generic relationships based on karyology) 497–510; *Minyichthys inusitatus* sp. nov. 774–776; *Niphon spinosus* (intrafamilial relationships) 777–787; *Liolaemus fabiani* sp. nov. 788–790; *Liophis (amazonicus, chrysostomus, miliaris, mossoroensis, purpurans)* (taxon. status) 791–802; *Litoria umbonata* sp. nov. 803–808; *Poecilia formosa* (lectotype designation) 816–817; *Poecilia (Mollienesia)* spp. (checklist & key to spp.) 817–822; *Minascaecilia sartorio* gen. & sp. nov. 857–863; *Modicus* gen. nov. (*minimus* sp. nov., *tangaroa* sp. nov.) 863–868; *Gastrotheca (bufona, cornuta, dendronastes* sp. nov.) (new sp., taxon. of trans-Andean marsupial frogs) 868–874; *Wetmorella (albofasciata, nigropinnata)* (revision, key to spp.) 875–883; *Pseudoeurycea (altamontana, leprosa, longicauda* sp. nov., *robertsi)* (new sp., relationships, electrophoretic & morphometric analyses) 884–894; *Rana (Paa) (annandalii, blanfordii, polunini, rara* sp. nov., *rostandi)* 895–901; Ostraciidae (aracanin genera (phylogenetic relationships of) 902–917; cryptodiran turtles (classif. & phylogeny of higher categories) 918–932; *Etheostoma (euzonum euzonum, euzonum erizonum)* (subspeciation, relationships) 1042–1050; *Thorius (dubitus, macdougalli, minutissimus, narisovalis, pennatulus narismagnus, p. pennatulus, pulmonaris, troglodytes,* spp. undescr.) (genetic relationships, taxon. identity, electrophoretic analyses) 1051–1073; *Boleosoma phlox, Ioa vigil, Etheostoma spectabile pulchellum, Percina shumardi* (synon.) 1082–1083; *Coryphaenoides (alateralis, thelestomus)* (validity of spp.) 1083–1086.

TAIL AUTOTOMY, *Podarcis taurica ionica* 991–1001.
TECHNIQUES, *Chrysemys picta belli, Pseudemys scripta elegans* (blood collecting by cardiac puncture) 522–523; resin-coated fish models for behavior studies 547–549; silver-staining of spermatozoa 974–978; stat. analysis on food and growth relationships 1090–1092.
TEMPERATURE, *Sceloporus variabilis* (testicular pigmentation & thermal physiol. of testis) 155–161; *Micropterus salmoides* (evol. in post-thermal environ.) 193–197; *Chrysemys picta* (as cue for seasonal gonadal cycle) 347–358; *Ambystoma* (*jeffersonianum, maculatum*) (preferred & acclimation temps.) 1001–1005.

VARIATION, *Fundulus zebrinus* (meristic & morphom. clinal variation) 55–67; *Rana clamitans* (size-related variation in calls) 141–155; *Gambusia affinis* (environ. & allele variation) 182–193; *Hyla gratiosa* (variation in growth & survival of larvae) 232–237; *Poeciliopsis lucida* (meristic variation) 287–299; *Acris crepitans* (temporal & geogr. variation in color morph proportions) 300–311; *Oryzias latipes* (allozymic variation in popns.) 311–318; *Anolis carolinensis* (genetic variation & popn.) 523–529; *Thamnophis* (*canascens, melanogaster*) (color & pattern variation) 530–534; *Lucania* (*goodei, interioris, parva*) (electrophoretic variation) 564–570; *Ambystoma t. tigrinum* (in body size & juv. production) 608–616; *Notophthalmus viridescens dorsalis* (variation in popn. density; effect on growth, survival & production) 629–639; neotropical hylid frogs (acoustic variation in 3 communities) 639–649; *Salmo gairdneri* (chromosomal variation) 650–662; *Dicamptodon* (*aterrimus, copei, ensatus*) (genetic variation) 679–691; *Fundulus heteroclitus* (of egg morphol. with geogr.) 726–740; *Opheodrys aestivus* (in stored lipids & reprod.) 741–745; *Liophis* (*amazonicus, chrysostomus, miliaris, mossoroensis, purpurans*) (geogr. variation) 791–802; *Pseudoeurycea* (*altamontana, leprosa, longicauda* sp. nov., *robertsi*) (intraspecific variation in *P. leprosa*) 884–894; *Gambusia affinis* (variation in life history traits among experimental popns.) 953–963; *Rana* (*clamitans, pipiens, sylvatica*), *Ptychadena mascareniensis* (dark-pigment variants) 979–990; *Desmognathus f. fuscus* (microgeographic size variation) 1101–1104; *Etheostoma* (*euzonum erizonum, e. euzonum*) (in characters with locality) 1042–1050.

ZOOGEOGRAPHY, *Chelonia mydas* (popns. with divergent nesting locations but shared feeding ranges) 121–125; *Anolis carolinensis* (historical biogeogr.) 523–529; *Liophis* (*amazonicus, chrysostomus, miliaris, mossoroensis, purpurans*) 791–802; "*Thalassoma*" larvae (occurrence in E. Pacific Barrier) 826–828.

Copeia

1983

Published by
THE AMERICAN SOCIETY OF ICHTHYOLOGISTS
AND HERPETOLOGISTS

CONTENTS—1983, Nos. 1–4

ABLE, KENNETH W. (*see* Morin, Richard P.)
ADAMS, M. (*see* Odendaal, F. J.)
ALEVIZON, WILLIAM (*see* Freeman, Steven)
ALLEN, LARRY G. AND JANET M. ILG. Larval development of the northern clingfish, *Gobiesox maeandricus* 551
ALLENDORF, FRED W. (*see* Daugherty, Charles H.) (*see* Phelps, Stevan R.)
ALLISON, IRA S. AND CARL E. BOND. Identity and probable age of salmonids from surface deposits at Fossil Lake, Oregon 563
ANDREWS, ROBIN M. AND A. STANLEY RAND. Limited dispersal of juvenile *Anolis limifrons* 429
ANGUS, ROBERT A. AND R. JACK SCHULTZ. Meristic variation in homozygous and heterozygous fish 287
AOKI, RIOSUKE. A new generic allocation of *Tomistoma machikanense*, a fossil crocodilian from the Pleistocene of Japan 89
ARNOLD, STEVAN J. (*see* Garland, Theodore, Jr.)
AYLING, A. M. AND JOHN R. PAXTON. *Odax cyanoallix*, a new species of odacid fish from northern New Zealand 95
BAILEY, JACK E., BRUCE L. WING AND JOYCE H. LANDINGHAM. Juvenile Pacific sandfish, *Trichodon trichodon*, associated with pink salmon, *Oncorhynchus gorbuscha*, fry in the nearshore area, southeastern Alaska 549
BAIRD, TROY A. Influence of social and predatory stimuli on the air-breathing behavior of the African clawed frog, *Xenopus laevis* 411
BALLINGER, ROYCE E. AND VICKY L. HOLSCHER. Assimilation efficiency and nutritive state in the striped plateau lizard, *Sceloporus virgatus* (Sauria: Iguanidae) 838
BALSANO, JOSEPH S. (*see* Monaco, Paul J.)
BAUER, BRUCE H., GERRY R. DINKINS AND DAVID A. ETNIER. Discovery of *Noturus baileyi* and *N. flavipinnis* in Citico Creek, Little Tennessee River system 558
BAUER, BRUCE H. (*see* Burkhead, Noel M.) (*see* Timmons, Tom J.)
BEAMISH, F. W. H. AND E. J. THOMAS. Potential and actual fecundity of the "paired" lampreys, *Ichthyomyzon gagei* and *I. castaneus* 367
BECKWITT, RICHARD. Genetic structure of *Genyonemus lineatus, Seriphus politus* (Sciaenidae) and *Paralabrax clathratus* (Serranidae) in southern California 691
BICKHAM, JOHN W. AND JOHN L. CARR. Taxonomy and phylogeny of the higher categories of cryptodiran turtles based on a cladistic analysis of chromosomal data 918
BOND, CARL E. (*see* Allison, Ira S.)
BORG, BERTIL AND ROBERT VAN DEN HURK. Oocytes in the testes of the three-spined stickleback, *Gasterosteus aculeatus* 259
BRANCH, LYN C. Social behavior of the tadpoles of *Phyllomedusa vaillanti* 420
BRINTON, STEPHEN L. (*see* Greenfield, David W.)
BRODIE, EDMUND D., JR. (*see* Ducey, Peter K.)
BRUNETTI, VALERIE (*see* Kallman, Klaus D.)
BRUSCA, RICHARD C. AND MATTHEW R. GILLIGAN. Tongue replacement in a marine fish (*Lutjanus guttatus*) by a parasitic isopod (Crustacea: Isopoda) 813
BUCCI-INNOCENTI, STEFANIA, MATILDE RAGGHIANTI AND GIORGIO MANCINO. Investigations of karyology and hybrids in *Triturus boscai* and *T. vittatus*, with a reinterpretation of the species groups within *Triturus* (Caudata: Salamandridae) 662
BULL, C. M. (*see* Odendaal, F. J.)
BURKHEAD, NOEL M. AND BRUCE H. BAUER. An intergeneric cyprinid hybrid, *Hybopsis monacha* × *Notropis galacturus*, from the Tennessee River drainage 1074
BURR, BROOKS M. AND ROBERT C. CASHNER. *Campostoma pauciradii*, a new cyprinid fish from southeastern United States, with a review of related forms 101
BUTTS, DAVID E. (*see* Howell, W. Mike)
CAMPBELL, JONATHAN A. (*see* Wake, Marvalee H.)
CARUSO, JOHN H. The systematics and distribution of the lophiid anglerfishes: II. Revisions of the genera *Lophiomus* and *Lophius* 11
CARR, JOHN L. (*see* Bickham, John W.)
CASHNER, ROBERT C. (*see* Burr, Brooks M.)
CHESSER, RONALD K. (*see* Smith, Michael W.)
CHISZAR, DAVID (*see* Scudder, Kent M.)
CHONDROPOULOS, B. P. AND J. J. LYKAKIS. Ecology of the Balkan wall lizard, *Podarcis taurica ionica* (Sauria: Lacertidae) from Greece 991

Clavijo, Ileana E. Pair spawning and formation of a lek-like mating system in the parrotfish *Scarus vetula* 253
Clements, William H. and Robert J. Livingston. Overlap and pollution-induced variability in the feeding habits of filefish (Pisces: Monacanthidae) from Apalachee Bay, Florida 331
Cole, Kathleen S. Protogynous hermaphroditism in a temperate zone territorial marine goby, *Coryphopterus nicholsi* 809
Coleman, Felicia. *Hemanthias peruanus*, another hermaphroditic anthiine serranid 252
Congdon, Justin D., Donald W. Tinkle and Philip C. Rosen. Egg components and utilization during development in aquatic turtles 264
Cordes, Lynn E. (*see* Page, Lawrence M.)
Crabtree, Roy E. Confirmation of the validity of *Coryphaenoides alateralis* as distinct from *Coryphaenoides thelestomus* based on new captures from the North Atlantic 1083
Creekmore, Jay S. (*see* Stephens, Gregory A.)
Crews, David (*see* McKenzie, William D., Jr.)
Critchley, K. (*see* Wood, J. R.)
Csányi, Vilmos (*see* Nagy, András)
Cuthbert, Glen J. (*see* Stevens, John D.)
Daniel, Richard E. (*see* Perrill, Stephen A.)
Daugherty, Charles H., Fred W. Allendorf, William W. Dunlap and Kathy L. Knudsen. Systematic implications of geographic patterns of genetic variation in the genus *Dicamptodon* 679
Davic, Robert D. Microgeographic body size variation in *Desmognathus fuscus fuscus* salamanders from western Pennsylvania 1101
Davies, Margaret (*see* Tyler, Michael J.)
Davis, John and R. Glenn Ford. Home range in the western fence lizard (*Sceloporus occidentalis occidentalis*) 933
Dawson, C. E. Western Atlantic occurrence of the genus *Minyichthys* (Pisces: Syngnathidae), with description of *M. inusitatus* n. sp. 774
Diaz, Nelson F., Jose Valencia and Michel Sallaberry. Life history and phylogenetic relationships of *Insuetophrynus acarpicus* (Anura: Leptodactylidae) 30
DiMichele, Leonard (*see* Taylor, Malcolm H.)
Dinkins, Gerry R. (*see* Bauer, Bruce H.)
Ditty, James G. and Frank M. Truesdale. Comparative larval development of *Peprilus burti, P. triacanthus* and *P. paru* (Pisces: Stromateidae) from the western North Atlantic 397
Dixon, James R. Taxonomic status of the South American snakes *Liophis miliaris, L. amazonicus, L. chrysostomus, L. mossoroensis* and *L. purpurans* (Colubridae: Serpentes) 791
Dominey, Wallace J. Mobbing in colonially nesting fishes, especially the bluegill, *Lepomis macrochirus* 1086
Drewry, George E. and A. Stanley Rand. Characteristics of an acoustic community: Puerto Rican frogs of the genus *Eleutherodactylus* 941
Drummond, Hugh. Adaptiveness of island nest-sites of green iguanas and slider turtles 529
Dubois, Alain and Masafumi Matsui. A new species of frog (genus *Rana*, subgenus *Paa*) from western Nepal (Amphibia: Anura) 895
Ducey, Peter K. and Edmund D. Brodie, Jr. Salamanders respond selectively to contacts with snakes: survival advantage of alternative antipredator strategies 1036
Duellman, William E. A new species of marsupial frog (Hylidae: *Gastrotheca*) from Colombia and Ecuador 868
Duellman, William E. and Rebecca A. Pyles. Acoustic resource partitioning in anuran communities 639
Dugan, Beverly (*see* Rand, A. Stanley)
Duggins, C. F., Jr., A. A. Karlin and K. G. Relyea. Electrophoretic variation in the killifish genus *Lucania* 564
Dunham, Arthur E. (*see* Tinkle, Donald W.)
Dunlap, William W. (*see* Daugherty, Charles H.)
Echternacht, Arthur C. (*see* Wade, Julia K.)
Egami, Nobuo (*see* Sakaizumi, Mitsuru)
Etheridge, Kay, Lawrence C. Wit and Jeffrey C. Sellers. Hibernation in the lizard *Cnemidophorus sexlineatus* (Lacertilia: Teiidae) 206
Etnier, David A. (*see* Bauer, Bruce H.)
Evans, Susan E. Mandibular fracture and inferred behavior in a fossil reptile 845
Fisher, Robert A. Protandric sex reversal in *Gonostoma elongatum* (Pisces: Gonostomatidae) from the eastern Gulf of Mexico 554

FITZSIMONS, JOHN MICHAEL (*see* Uyeno, Teruya)
FLATER, GARY (*see* Guillette, Louis J., Jr.)
FLOYD, HENRY B. AND THOMAS A. JENSSEN. Food habits of the Jamaican lizard *Anolis opalinus*: resource partitioning and seasonal effects examined 319
FORD, R. GLENN (*see* Davis, John)
FORESTER, DON C. Duration of the brooding period in the mountain dusky salamander (*Desmognathus ochrophaeus*) and its influence on aggression toward conspecifics 1098
FORMAS, J. R. AND M. INES VERA. Karyological relationships among frogs of the genus *Alsodes*, with description of the karyotypes of *A. vanzolinii* and *A. verrucosus* 1104
FRAZER, NAT B. Effect of tidal cycles on loggerhead sea turtles (*Caretta caretta*) emerging from the sea 516
FREEMAN, STEVEN AND WILLIAM ALEVIZON. Aspects of territorial behavior and habitat distribution of the fairy basslet *Gramma loreto* 829
GANZHORN, DARRIE AND PAUL LICHT. Regulation of seasonal gonadal cycles by temperature in the painted turtle, *Chrysemys picta* 347
GARLAND, THEODORE, JR. AND STEVAN J. ARNOLD. Effects of a full stomach on locomotory performance of juvenile garter snakes (*Thamnophis elegans*) 1092
GARTNER, JOHN V., JR. Sexual dimorphism in the bathypelagic gulper eel *Eurypharynx pelecanoides* (Lyomeri: Eurypharyngidae), with comments on reproductive strategy 560
GATES, J. EDWARD (*see* Stauffer, Jay R., Jr.)
GATTEN, ROBERT E., JR. (*see* Stockard, Matthew E.)
GEDDES, CYNTHIA (*see* Ruben, John A.)
GERLACH, JANE M. Characters for distinguishing larvae of carp, *Cyprinus carpio*, and goldfish, *Carassius auratus* 116
GILLIGAN, MATTHEW R. (*see* Brusca, Richard C.)
GILMORE, R. GRANT. Observations on the embryos of the longfin mako, *Isurus paucus*, and the bigeye thresher, *Alopias superciliosus* 375
GLUESING, ERNEST A. Collared lizard predation: the effects of conspicuous morphology and movement 835
GOLDBERG, STEPHEN R. (*see* Vitt, Laurie J.)
GOODFELLOW, WILLIAM L. (*see* Stauffer, Jay R., Jr.)
GOW, GRAEME (*see* King, Max)
GRAY, ROBERT H. Seasonal, annual and geographic variation in color morph frequencies of the cricket frog, *Acris crepitans*, in Illinois 300
GREELEY, MARK S., JR. AND ROBERT MACGREGOR, III. Annual and semilunar reproductive cycles of the Gulf killifish, *Fundulus grandis*, on the Alabama Gulf Coast 711
GREENFIELD, DAVID W. *Gambusia xanthosoma*, a new species of poeciliid fish from Grand Cayman Island, BWI 457
GREENFIELD, DAVID W., TERESA A. GREENFIELD AND STEPHEN L. BRINTON. Spatial and trophic interactions between *Gambusia sexradiata* and *Gambusia puncticulata yucatana* (Pisces: Poeciliidae) in Belize, Central America 598
GREENFIELD, TERESA A. (*see* Greenfield, David W.)
GREGORY, LINDA A. (*see* Gregory, Patrick T.)
GREGORY, PATRICK T., LINDA A. GREGORY AND J. MALCOLM MACARTNEY. Color-pattern variation in *Thamnophis melanogaster* 530
GROUT, DOUGLAS E. A case of hermaphroditism in the rainbow smelt, *Osmerus mordax* 812
GUILLETTE, LOUIS J., JR., JOHN WEIGEL AND GARY FLATER. Unilateral testicular pigmentation in the Mexican lizard *Sceloporus variabilis* 155
HAIRSTON, NELSON G. Growth, survival and reproduction of *Plethodon jordani*: trade-offs between selective pressures 1024
HALL, JOHN G. (*see* Smith, G. R.)
HANKEN, JAMES. Genetic variation in a dwarfed lineage, the Mexican salamander genus *Thorius* (Amphibia: Plethodontidae): taxonomic, ecologic and evolutionary implications 1051
HARDY, GRAHAM S. A new genus and two new species of clingfishes (Gobiesocidae) from New Zealand 863
HELFMAN, GENE S. Resin-coated fishes: a simple model technique for in situ studies of fish behavior 547
HILLMAN, STANLEY S. AND MALCOLM S. LEA. Aerial activity oxygen consumption during metamorphosis of the bullfrog, *Rana catesbeiana* 407
HINCHLIFFE, P. M. (*see* Richardson, K. C.)
HIRSCH, KARL F. Contemporary and fossil chelonian eggshells 382
HOLSCHER, VICKY L. (*see* Ballinger, Royce E.)

HOWELL, W. MIKE AND DAVID E. BUTTS. Silver staining of spermatozoa from living and preserved museum fishes: a new taxonomic approach 974
HUBBS, CLARK (*see* Miller, Robert Rush)
HUNT, LAWRENCE E. A nomenclatural rearrangement of the genus *Anniella* (Sauria: Anniellidae) 79
HURK, ROBERT VAN DEN (*see* Borg, Bertil)
HURST, CHARLES J. (*see* Ramer, John D.)
ILG, JANET M. (*see* Allen, Larry G.)
INNES, DAVID J. (*see* Smith, G. R.)
JACOBSON, LARRY (*see* Pierce, Benjamin A.)
JAMES, ANTHONY H. (*see* James, David K.)
JAMES, DAVID K., LEWIS PETRINOVICH, THOMAS L. PATTERSON AND ANTHONY H. JAMES. Predation of white-crowned sparrow nestlings by the western terrestrial garter snake in San Francisco, California 511
JANSEN, DAVID W. A possible function of the secretion of Duvernoy's gland 262
JENKINS, ROBERT L. Observations on the commensal relationship of *Nomeus gronovii* with *Physalia physalis* 250
JENSEN, GARY L. AND WILLIAM L. SHELTON. Gonadal differentiation in relation to sex control of grass carp, *Ctenopharyngodon idella* (Pisces: Cyprinidae) 749
JENSSEN, THOMAS A. (*see* Floyd, Henry B.) (*see* Ramer, John D.)
JOHNSON, G. DAVID. *Niphon spinosus*: a primitive epinepheline serranid, with comments on the monophyly and intrarelationships of the Serranidae 777
JONES, JEFFERSON C. (*see* Smith, Michael H.)
KALLMAN, KLAUS D. The sex determining mechanism of the poeciliid fish, *Xiphophorus montezumae*, and the genetic control of the sexual maturation process and adult size 775
KALLMAN, KLAUS D. AND VALERIE BRUNETTI. Genetic basis of three mutant color varieties of *Xiphophorus maculatus*: the gray, gold and ghost platyfish 170
KALLMAN, KLAUS D. (*see* McKenzie, William D., Jr.)
KARLIN, A. A. (*see* Duggins, C. F., Jr.)
KING, MAX AND GRAEME GOW. A new species of *Oedura* (Gekkonidae: Reptilia) from the Alligator Rivers region of northern Australia 445
KLUGE, ARNOLD G. Cladistic relationships among gekkonid lizards 465
KNUDSEN, KATHY L. (*see* Daugherty, Charles H.)
KOBAYASHI, DANI, WILLIAM J. MAUTZ AND KENNETH A. NAGY. Evaporative water loss: humidity acclimation in *Anolis carolinensis* lizards 701
KOCHER, THOMAS D. AND KENNETH R. MCKAYE. Defense of heterospecific cichlids by *Cyrtocara moorii* in Lake Malawi, Africa 544
KOEHN, RICHARD K. (*see* Smith, G. R.)
KOFRON, CHRISTOPHER P. Female reproductive cycle of the neotropical snail-eating snake *Sibon sanniola* in northern Yucatan, Mexico 963
KREKORIAN, C. O'NEIL. Population density of the desert iguana, *Dipsosaurus dorsalis* (Reptilia: Iguanidae), in southern California 268
LANDINGHAM, JOYCE H. (*see* Bailey, Jack E.)
LEA, MALCOLM S. (*see* Hillman, Stanley S.)
LEIS, JEFFREY M. Coral reef fish larvae (Labridae) in the East Pacific Barrier 826
LICHT, PAUL (*see* Ganzhorn, Darrie)
LIU, EDWIN H. (*see* Smith, Michael H.)
LIVINGSTON, ROBERT J. (*see* Clements, William H.)
LUCAS, SPENCER G. (*see* Storrs, Glenn W.)
LYKAKIS, J. J. (*see* Chondropoulos, B. P.)
LYNCH, JAMES F., DAVID B. WAKE AND SUH Y. YANG. Genic and morphological differentiation in Mexican *Pseudoeurycea* (Caudata: Plethodontidae), with a description of a new species 884
MACARTNEY, J. MALCOLM (*see* Gregory, Patrick T.)
MACGREGOR, ROBERT, III (*see* Greeley, Mark S., Jr.)
MANCINO, GIORGIO (*see* Bucci-Innocenti, Stefania)
MATSUI, MASAFUMI (*see* Dubois, Alain)
MAUTZ, WILLIAM J. (*see* Kobayashi, Dani)
MAYHEW, DAVID A. A new hybrid cross, *Notropis atherinoides* × *Notropis volucellus* (Pisces: Cyprinidae), from the lower Monongahela River, western Pennsylvania 1077
MCCRACKEN, GARY F. (*see* Wade, Julia K.)

McElman, James F. Comparative embryonic ecomorphology and the reproductive guild classification of walleye, *Stizostedion vitreum,* white sucker, *Catostomus commersoni* 246
McKaye, Kenneth R. (*see* Kocher, Thomas D.)
McKenzie, William D., Jr., David Crews, Klaus D. Kallman, David Policansky and Joel J. Sohn. Age, weight and the genetics of sexual maturation in the platyfish, *Xiphophorus maculatus* 770
Mendonça, Mary T. Movements and feeding ecology of immature green turtles (*Chelonia mydas*) in a Florida lagoon 1013
Metter, Dean E. (*see* Peterson, Chris L.)
Miller, Robert Rush. Checklist and key to the mollies of Mexico (Pisces: Poeciliidae: *Poecilia,* subgenus *Mollienesia*) 817
Miller, Robert Rush and Clark Hubbs. Designation of a lectotype for the Amazon molly, *Poecilia formosa* (Pisces: Poeciliidae) 816
Miller, Robert Rush (*see* Poss, Stuart G.) (*see* Uyeno, Teruya)
Mitton, Jeffry B. (*see* Pierce, Benjamin A.)
Moll, Don. A proposed origin of nesting location divergence and a shared feeding range in Brazilian green turtle populations 121
Monaco, Paul J., Ellen M. Rasch and Joseph S. Balsano. The occurrence of superfetation in the Amazon molly, *Poecilia formosa,* and its related sexual species 969
Monostory, Zsuzsa (*see* Nagy, András)
Montanucci, Richard R. Natural hybridization between two species of collared lizards (*Crotaphytus*) 1
Morin, Peter J. Competitive and predatory interactions in natural and experimental populations of *Notophthalmus viridescens dorsalis* and *Ambystoma tigrinum* 628
Morin, Richard P. and Kenneth W. Able. Patterns of geographic variation in the egg morphology of the fundulid fish, *Fundulus heteroclitus* 726
Moriwaki, Kazuo (*see* Sakaizumi, Mitsuru)
Nace, George W. (*see* Richards, Christina M.)
Nagy, András, Zsuzsa Monostory and Vilmos Csányi. Rapid development of the clonal state in successive gynogenetic generations of carp (*Cyprinus carpio*) 745
Nagy, Kenneth A. (*see* Kobayashi, Dani)
Narins, Peter M. Divergence of acoustic communication systems of two sibling species of eleutherodactylid frogs 1089
Nelson, Gareth. *Anchoa argentivittata,* with notes on other eastern Pacific anchovies and the Indo-Pacific genus *Encrasicholina* 48
Núñez, Herman and José Yáñez. *Ctenoblepharis adituvelatus* new species, a lizard from northern Chile (Reptilia: Iguanidae) 454
Núñez, Herman (*see* Yáñez, José L.)
Odendaal, F. J., C. M. Bull and M. Adams. Genetic divergence between two morphologically similar *Ranidella* species (Anura: Leptodactylidae) 275
Odendaal, F. J., C. M. Bull and S. R. Telford. The vocabulary of calls of *Ranidella riparia* (Anura: Leptodactylidae) 534
Page, Lawrence M. Identification of the percids, *Boleosoma phlox* Cope and *Ioa vigil* Hay 1082
Page, Lawrence M. and Lynn E. Cordes. Variation and systematics of *Etheostoma euzonum,* the Arkansas saddled darter (Pisces: Percidae) 1042
Parmenter, C. J. Reproductive migration in the hawksbill turtle (*Eretmochelys imbricata*) 271
Patterson, Thomas L. (*see* James, David K.)
Paxton, John R. (*see* Ayling, A. M.)
Perrill, Stephen A. and Richard E. Daniel. Multiple egg clutches in *Hyla regilla, H. cinerea* and *H. gratiosa* 513
Peterson, Chris L., Robert F. Wilkinson, Jr., Milton S. Topping and Dean E. Metter. Age and growth of the Ozark hellbender (*Cryptobranchus alleganiensis bishopi*) 225
Petranka, James W. Fish predation: a factor affecting the spatial distribution of a stream-breeding salamander 624
Petrinovich, Lewis (*see* James, David K.)
Phelps, Stevan R. and Fred W. Allendorf. Genetic identity of pallid and shovelnose sturgeon (*Scaphirhynchus albus* and *S. platorynchus*) 696
Pierce, Benjamin A., Jeffry B. Mitton, Larry Jacobson and Francis L. Rose. Head shape and size in cannibal and noncannibal larvae of the tiger salamander from west Texas 1006
Plummer, Michael V. Annual variation in stored lipids and reproduction in green snakes (*Opheodrys aestivus*) 741

POLICANSKY, DAVID (*see* McKenzie, William D., Jr.)
POSS, STUART G. AND ROBERT RUSH MILLER. Taxonomic status of the plains killifish, *Fundulus zebrinus* 55
PYLES, REBECCA A. (*see* Duellman, William E.)
RAGGHIANTI, MATILDE (*see* Bucci-Innocenti, Stefania)
RAJ, UDAY AND JOHNSON SEETO. A new species of *Paracaesio* (Pisces: Lutjanidae) from the Fiji Islands 450
RAMER, JOHN D., THOMAS A. JENSSEN AND CHARLES J. HURST. Size-related variation in the advertisement call of *Rana clamitans* (Anura: Ranidae), and its effect on conspecific males 141
RAMSEY, JOHN S. (*see* Timmons, Tom J.)
RAND, A. STANLEY AND BEVERLY DUGAN. Structure of complex iguana nests 705
RAND, A. STANLEY (*see* Andrews, Robin M.) (*see* Drewry, George E.)
RANDALL, JOHN E. Revision of the Indo-Pacific labrid fish genus *Wetmorella* 875
RASCH, ELLEN M. (*see* Monaco, Paul J.)
REWCASTLE, STEPHEN COMPTON. Fundamental adaptations in the lacertilian hind limb: a partial analysis of the sprawling limb posture and gait 476
RELYEA, K. G. (*see* Duggins, C. F., Jr.)
RICHARDS, CHRISTINA M. AND GEORGE W. NACE. Dark pigment variants in anurans: classification, new descriptions, color changes and inheritance 979
RICHARDSON, K. C. AND P. M. HINCHLIFFE. Caudal glands and their secretions in the western spiny-tailed gecko, *Diplodactylus spinigerus* 161
ROBERTS, TYSON R. Revision of the South and Southeast Asian sisorid catfish genus *Bagarius*, with description of a new species from the Mekong 435
ROSE, FRANCIS L. (*see* Pierce, Benjamin A.)
ROSEN, PHILIP C. (*see* Congdon, Justin D.)
ROWLAND, WILLIAM J. Interspecific aggression in sticklebacks: *Gasterosteus aculeatus* displaces *Apeltes quadracus* 541
RUBEN, JOHN A. AND CYNTHIA GEDDES. Some morphological correlates of striking in snakes 221
SAKAIZUMI, MITSURU, KAZUO MORIWAKI AND NOBUO EGAMI. Allozymic variation and regional differentiation in wild populations of the fish *Oryzias latipes* 311
SALDAÑA, JUAN AND BARNEY VENABLES. Energy compartmentalization in a migratory fish, *Prochilodus mariae* (Prochilodontidae), of the Orinoco River 617
SALLABERRY, MICHEL (*see* Diaz, Nelson F.)
SCHOCH, ROBERT M. (*see* Storrs, Glenn W.)
SCHULTZ, R. JACK (*see* Angus, Robert A.)
SCOTT, STEPHANIE L. (*see* Smith, Michael H.)
SCUDDER, KENT M., DAVID CHISZAR AND HOBART M. SMITH. Effect of environmental odors on strike-induced chemosensory searching by rattlesnakes 519
SEETO, JOHNSON (*see* Raj, Uday)
SEIGEL, RICHARD A. Natural survival of eggs and tadpoles of the wood frog, *Rana sylvatica* 1096
SELLERS, JEFFREY C. (*see* Etheridge, Kay)
SEMLITSCH, RAYMOND D. Structure and dynamics of two breeding populations of the eastern tiger salamander, *Ambystoma tigrinum* 608
SEMLITSCH, RAYMOND D. AND CAROLYN A. WEST. Aspects of the life history and ecology of Webster's salamander, *Plethodon websteri* 339
SHAFFER, H. BRADLEY. Biosystematics of *Ambystoma rosaceum* and *A. tigrinum* in northeastern Mexico 67
SHELDON, ANDREW L. Food and growth relationships: a comment on statistical analysis 1090
SHELTON, WILLIAM L. (*see* Jensen, Gary L.)
SHINE, RICHARD. Arboreality in snakes: ecology of the Australian elapid genus *Hoplocephalus* 198
SMITH, G. R., JOHN G. HALL, RICHARD K. KOEHN AND DAVID J. INNES. Taxonomic relationships of the Zuni mountain sucker, *Catostomus discobolus yarrowi* 37
SMITH, HOBART M. (*see* Scudder, Kent M.)
SMITH, J. D. (*see* Smith, R. J. F.)
SMITH, MICHAEL H., MICHAEL W. SMITH, STEPHANIE L. SCOTT, EDWIN H. LIU AND JEFFERSON C. JONES. Rapid evolution in a post-thermal environment 193
SMITH, MICHAEL H. (*see* Smith, Michael W.)
SMITH, MICHAEL W., MICHAEL H. SMITH AND RONALD K. CHESSER. Biochemical genetics of mosquitofish. I. Environmental correlates, and temporal and spatial heterogeneity of allele frequencies within a river drainage 182
SMITH, MICHAEL W. (*see* Smith, Michael H.)

SMITH, R. J. F. AND J. D. SMITH. Seasonal loss of alarm substance cells in *Chrosomus neogaeus, Notropis venustus* and *N. whipplei* 822
SOHN, JOEL J. (*see* McKenzie, William D., Jr.)
STAUFFER, JAY R., JR., J. EDWARD GATES AND WILLIAM L. GOODFELLOW. Preferred temperature of two sympatric *Ambystoma* larvae: a proximate factor in niche segregation? 1001
STEINWASCHER, KURT AND JOSEPH TRAVIS. Influence of food quality and quantity on early larval growth of two anurans 238
STEPHENS, GREGORY A. AND JAY S. CREEKMORE. Blood collection by cardiac puncture in conscious turtles 522
STEVENS, J. D. Observations on reproduction in the shortfin mako *Isurus oxyrinchus* 126
STEVENS, JOHN D. AND GLEN J. CUTHBERT. Observations on the identification and biology of *Hemigaleus* (Selachii: Carcharhinidae) from Australian waters 487
STOCKARD, MATTHEW E. AND ROBERT E. GATTEN, JR. Activity metabolism of painted turtles (*Chrysemys picta*) 214
STORRS, GLENN W., SPENCER G. LUCAS AND ROBERT M. SCHOCH. Endocranial cast of an early Paleocene crocodilian from the San Juan Basin, New Mexico 842
TAYLOR, JEFFREY N. Field observations on the reproductive ecology of three species of armored catfishes (Loricariidae: Loricariinae) in Paraguay 257
TAYLOR, MALCOLM H. AND LEONARD DIMICHELE. Spawning site utilization in a Delaware population of *Fundulus heteroclitus* (Pisces: Cyprinodontidae) 719
TELFORD, S. R. (*see* Odendaal, F. J.)
THOMAS, E. J. (*see* Beamish, F. W. H.)
THOMPSON, SUE. Homing in a territorial reef fish 832
THORGAARD, GARY H. Chromosomal differences among rainbow trout populations 650
TIMMONS, TOM J., JOHN S. RAMSEY AND BRUCE H. BAUER. Life history and habitat of the blackfin sucker, *Moxostoma atripinne* (Osteichthyes: Catostomidae) 538
TINKLE, DONALD W. AND ARTHUR E. DUNHAM. Demography of the tree lizard, *Urosaurus ornatus*, in central Arizona 585
TINKLE, DONALD W. (*see* Congdon, Justin D.)
TOPPING, MILTON S. (*see* Peterson, Chris L.)
TRAVIS, JOSEPH. Variation in growth and survival of *Hyla gratiosa* larvae in experimental enclosures 232
TRAVIS, JOSEPH (*see* Steinwascher, Kurt)
TRENDALL, JASPER T. Life history variation among experimental populations of the mosquitofish, *Gambusia affinis* 953
TRUESDALE, FRANK M. (*see* Ditty, James G.)
TYLER, JAMES C. (*see* Winterbottom, Richard)
TYLER, MICHAEL J. AND MARGARET DAVIES. A new species of *Litoria* (Anura: Hylidae) from Irian Jaya, New Guinea 803
UYENO, TERUYA, ROBERT RUSH MILLER AND JOHN MICHAEL FITZSIMONS. Karyology of the cyprinodontoid fishes of the Mexican family Goodeidae 497
VALENCIA, JOSE (*see* Diaz, Nelson F.)
VENABLES, BARNEY (*see* Saldaña, Juan)
VERA, M. INES (*see* Formas, J. R.)
VITT, LAURIE J. Reproduction and sexual dimorphism in the tropical teiid lizard *Cnemidophorus ocellifer* 359
VITT, LAURIE J. AND STEPHEN R. GOLDBERG. Reproductive ecology of two tropical iguanid lizards: *Tropidurus torquatus* and *Platynotus semitaeniatus* 131
WADE, JULIA K., ARTHUR C. ECHTERNACHT AND GARY F. MCCRACKEN. Genetic variation and similarity in *Anolis carolinensis* (Sauria: Iguanidae) 523
WAKE, DAVID B. (*see* Lynch, James F.)
WAKE, MARVALEE H. AND JONATHAN A. CAMPBELL. A new genus and species of caecilian from the Sierra de las Minas of Guatemala 857
WEIGEL, JOHN (*see* Guillette, Louis J., Jr.)
WEST, CAROLYN A. (*see* Semlitsch, Raymond D.)
WHITMORE, DONALD H. Introgressive hybridization of smallmouth (*Micropterus dolomieui*) and Guadalupe bass (*M. treculi*) 672
WILEY, JOHN E. Chromosome polymorphism in *Hyla chrysoscelis* 273
WILKINSON, ROBERT F., JR. (*see* Peterson, Chris L.)
WING, BRUCE L. (*see* Bailey, Jack E.)

WINTERBOTTOM, RICHARD AND JAMES C. TYLER. Phylogenetic relationships of aracanin genera of boxfishes (Ostraciidae: Tetraodontiformes) 902
WIT, LAWRENCE C. (*see* Etheridge, Kay)
WOOD, F. E. (*see* Wood, J. R.)
WOOD, J. R., F. E. WOOD AND K. CRITCHLEY. Hybridization of *Chelonia mydas* and *Eretmochelys imbricata* 839
YÁÑEZ, JOSÉ L. AND HERMAN NÚÑEZ. *Liolaemus fabiani*, a new species of lizard from northern Chile (Reptilia: Iguanidae) 788
YÁÑEZ, JOSÉ (*see* Núñez, Herman)
YANG, SUH Y. (*see* Lynch, James F.)
ZIMMERER, EDMUND J. Effect of salinity on the size-hierarchy effect in *Poecilia latipinna*, *P. reticulata* and *Gambusia affinis* 243